

Population-based versus hospital-based controls: are they comparable?

Alberto Ruano-Ravina^{a,b,c} / Mónica Pérez-Ríos^{a,c,e} / Juan Miguel Barros-Dios^{a,c,d}

^aDepartment of Preventive Medicine and Public Health, University of Santiago de Compostela, Spain;

^bGalician Agency for Health Technology Assessment, Galician Health Service, Spain;

^cCIBER de Epidemiología y Salud Pública, CIBERESP, Spain; Department of Public Health, Galician Regional Authority, Spain;

^dPreventive Medicine Service, Santiago de Compostela University Teaching Hospital, Spain.

(Controles poblaciones frente a controles hospitalarios: ¿son comparables?)

Abstract

Objective: To compare whether there are differences among hospital and population controls.

Methods: Two case-control studies were conducted on lung cancer risk factors in the Santiago de Compostela Public Health District. Whereas one used randomly chosen census-based population controls, the other used consecutive hospital controls that went to the reference hospital for non-smoking-related trivial interventions. The differences were analyzed using logistic regression. The dependent variable was type of control (hospital or population).

Results: Hospital controls had a similar tobacco habit than population controls, but consumed more alcohol. For those consuming more than 50 ml daily, the risk of being a hospital control was 4.83 (95%CI: 2.55-9.14).

Conclusions: There may be some differences between hospital and population-based controls, which must be taken into account in the design of case-control studies. It is necessary to ascertain whether such differences are reproduced at other geographic locations and whether they can affect estimation of exposure-disease

Key words: Epidemiologic studies. Case-control studies. Retrospective studies. Selection bias. Tobacco. Alcohol.

Resumen

Objetivo: Comparar si hay diferencias entre los controles poblacionales y los hospitalarios.

Métodos: Se llevaron a cabo dos estudios de casos y controles sobre factores de riesgo de cáncer de pulmón en el Área Sanitaria de Santiago de Compostela. En uno de los estudios los controles fueron seleccionados aleatoriamente entre la población general, y en el otro los controles hospitalarios fueron incluidos de manera consecutiva entre los individuos que acudían al hospital por intervenciones quirúrgicas banales no relacionadas con el consumo de tabaco. Las diferencias fueron analizadas mediante regresión logística. La variable dependiente fue el tipo de control (hospitalario o poblacional).

Resultados: Los controles hospitalarios y los poblacionales tenían un hábito tabáquico similar, pero los controles hospitalarios consumían más alcohol. Para los que consumían más de 50 ml al día, el riesgo de ser un control hospitalario fue de 4,83 (intervalo de confianza del 95%: 2,55-9,14).

Conclusiones: Podría haber algunas diferencias entre los controles poblacionales y los hospitalarios que deberían tenerse en cuenta cuando se diseñe un estudio de casos y controles. Es necesario saber si esas diferencias son similares en otras áreas geográficas y si podrían afectar a la estimación de las medidas de efecto entre exposición y enfermedad.

Palabras clave: Estudios epidemiológicos. Estudios de casos y controles. Estudios retrospectivos. Sesgo de selección. Tabaco. Alcohol.

Background

Case-control studies are one of the most frequently used designs in epidemiology, having been defined as an efficient version of cohort studies¹. Part of the validity of case-control studies resides in appropriate selection of controls. There are two major groups of controls, viz., population and hospital, yet few studies have compared the two. It has been claimed that, if hospital controls are properly selected, there should be no important differences as against population controls in terms of the risks obtained for any given disease².

This study sought to compare the characteristics of the above two groups of controls (hospital and popu-

Correspondence: Alberto Ruano-Ravina.
Department of Preventive Medicine and Public Health,
School of Medicine. University of Santiago de Compostela.
San Francisco, s/n. 15782 Santiago de Compostela. Spain.
E-mail: alberto.ruano@usc.es

Received: July 18, 2007.

Accepted: January 9, 2008.

lation) recruited in the same public health district for two case-control studies on risk factors for lung cancer.

Methods

The two groups of controls were enrolled for two lung cancer studies. The objectives of the respective studies were: in one case, to analyze the effect of radon exposure³; and in the other, to analyze the effect of environmental and genetic risk factors of lung cancer⁴. In both studies, participation was restricted to subjects over 35 years and the sampling was performed based on sex frequency among cases.

Hospital-based controls

Controls attending the Santiago de Compostela Hospital Complex Preoperative Unit for trivial surgery were recruited between May 1999-May 2000 through consecutive sampling. Subjects with previous neoplasms were excluded. The breakdown of these surgical interventions was as follows: 31%, cataracts; 30%, inguinal hernias; 11%, orthopedic surgery; and the remainder, non-smoking-related interventions. All interviews were conducted by a single interviewer at the hospital. Participation was 100%.

Population controls

A total of 500 subjects from the Health Area of the Santiago de Compostela Hospital Complex were initially selected, using the 1991 census. A proportional stratified random sampling was carried out, based on distribution by sex among cases and the population weighting of different sub-areas.

The following exclusion criteria were established for controls: less than 5 years uninterrupted residence in the current home and/or residence of any length in any dwelling that had undergone major structural alterations. Individuals suffering respiratory tract diseases or having a clinical history of any type of neoplasm, past or present, were excluded. Interviews were conducted at home by two interviewers. A total of 391 subjects met the eligibility criteria, 241 of whom finally participated as population controls, entailing a participation of 61.6%.

Data-source and definition of variables

The questionnaires were very similar in both studies, since they sought to measure the same effect (lung cancer) and a series of variables common to both studies

(tobacco use, alcohol, occupation, and diet). Both questionnaires had been previously tested on a group of persons who had similar characteristics, and all questions poorly drafted or ambiguous were duly amended or eliminated.

A smoker was defined as any subject who, at some stage in his/her life, had smoked a minimum of one cigarette per day for at least one year. An ex-smoker was defined as any subject who, at the date of the interview, had refrained from smoking for at least one year. Weekly alcohol consumption (in grams) was deemed to be the combined intake of different alcoholic beverages, such as wine, beer, and spirits. In addition, educational level was classified into no formal education, primary education, and secondary or higher education. We gathered information on individuals' occupations, and considered those defined as risk occupations for lung cancer⁵. Subjects were classified dichotomously as exposed or non-exposed, according to whether they had worked for a minimum of one year in an occupation deemed to be a risk for lung cancer.

Statistical analysis

Data were analyzed using unconditional logistic regression, calculating the probability of being a hospital control versus a population control. We considered the following independent variables: duration of smoking (classified into non-smokers, 1-30 years, and over 30 years); number of cigarettes smoked per day (classified into non-smokers, 1-20, and over 20), number of cigarette packs smoked over a lifetime (with non-smokers and smokers divided into equal parts, equivalent to moderate and heavy smokers); educational level (no formal education, primary education, and secondary or higher education); occupation and daily alcohol consumption (non-drinkers, moderate drinkers [1 to 50 ml per day] and heavy drinkers [> 50 ml per day]).

Practice application

A description of the study sample is shown in table 1. A total of 187 hospital and 241 population controls were included. There was a difference of six years in the mean age of the two groups, though the age range was similar. The percentage of individuals with secondary or higher education was likewise similar, but the percentage of individuals without any formal education was greater among hospital controls. Whereas the percentage of smokers was slightly higher among hospital controls, the number of cigarettes smoked per day and percentage of never-smokers was very much alike in both control

Table 1. Description of the sample of hospital and population controls

	Hospital controls (n = 187)	Population controls (n = 241)
Sex		
Males (%)	164 (87.7)	219 (90.9)
Age (years)		
Mean (SD)	62.5 (9.9)	56.9 (12.5)
Range	40-86	35-93
Studies		
No formal education (%)	55 (29.4)	22 (9.1)
Elementary education (%)	114 (61.0)	195 (80.9)
Secondary education or higher (%)	18 (9.6)	24 (10.0)
Tobacco consumption		
Current smokers ^a (%)	70 (38.1)	86 (35.7)
Mean cigarettes/day (SD)	13.0 (15.4)	11.6 (14.2)
Never smokers (%)	77 (41.2)	106 (44.0)
Alcohol consumption		
Mean weekly intake in ml (SD)	384.9 (371.4)	183.9 (192.3)
Non-drinkers (%)	29 (15.5)	79 (32.8)
Heavy drinkers (%) (> 50 ml per week)	80 (42.8)	80 (33.2)
Occupation		
Have not worked in risk occupations for lung cancer (%)	138 (64.5)	163 (67.7)
Have worked more than 1 year in risk occupations for lung cancer (%)	49 (35.5)	78 (32.3)

^aDaily smokers and occasional smokers.

groups. Alcohol consumption was higher among hospital controls, as was the percentage of drinkers.

Table 2 shows the probability of being a hospital control versus a population control. No important differences were observed between both in smoking habit and occupational exposure to carcinogens. Important differences appeared for alcohol consumption. The risk of being a hospital control was 2.76-fold higher for moderate drinkers, rising to 4.83-fold for heavy drinkers, which means that the higher the alcohol consumption, the higher the likelihood of being selected as a hospital control.

Few studies have compared directly hospital and population controls, despite the amount of theoretical literature on the topic. These results indicate that both control types are comparable in terms of factors viewed as potential confounding variables in many epidemiologic studies, e.g., educational level, number of cigarettes smoked per day, duration of smoking habit, and even risk occupations for lung cancer. The results also indicate, however, that there are important differences between hospital and population controls regarding alcohol consumption, with hospital controls having a greater likelihood of being drinkers.

The use of population controls is acknowledged as being more appropriate than that of hospital controls for

reasons of representativeness. Nevertheless, both have their advantages and their drawbacks. In the study region, public coverage covers approximately 95% of the population. The population controls were randomly selected on the basis of the census. There may be certain differential characteristics between the 5% that makes no use of public health and the remaining 95%, in aspects such as economic level (educational level), tobacco use (the lower classes tend to smoke more than the higher classes), and perhaps employment in occupations posing a risk for lung cancer.

The basic problem of using hospital or population controls is that of comparability. In the present study, both control types were comparable except in alcohol consumption (and educational level). It might be thought that this is due to the medical profiles in respect of which hospital controls were selected. However, none of these disorders (inguinal hernia, cataracts and orthopedic surgery in over 95% of controls) is, in principle, associated with alcohol consumption, so that they could not account for the differences observed. Another possible explanation might be non-response or participation bias. In general, participation by hospital controls tends to be higher than that of population controls. Hospital setting may increase participation, while the fact of being a healthy (and in many cases, a working) person reduces participation among population controls. Participation among hospital controls was 100%, whereas it was only 61.6% of the population controls. It may be that a higher proportion of heavier drinkers decline to participate among population than among hospital controls, which could account for the results obtained. Insofar as memory bias is concerned, hospital controls are reported to have as much time as cases do to think about the causes of their disease, thus preventing any possible differential information bias⁶. Drinkers may be more reluctant to participate when they are approached in their home environment or because they are more difficult to locate in this setting.

Our study possesses a series of advantages; the two studies from which the controls were taken were designed to study the same disease. Consequently, the questionnaires used were essentially similar. Accordingly, it can be said that, in this particular instance of a case-control study—one in which the «cases» were hospital controls—the principle of comparable measures has been fulfilled⁷. A further advantage resides in the fact that, from the very outset, both studies were undertaken in the same public health district, a factor that would, a priori, make both groups of controls highly comparable⁷. This means that the distribution of the different study exposures (and of other variables that have not been evaluated) is exactly the same in the population base from which the two groups of controls were drawn. Both studies used the same inclusion and exclusion criteria (save

Table 2. Probability of being a hospital control

	Hospital controls	Population controls	OR ^a	95%CI	OR ^b	95%CI
Education						
No formal education	55	22	1	–	1	–
Primary education	114	195	0.29	0.16-0.51	0.32	0.18-0.57
Secondary school or higher	18	18	0.46	0.20-1.04	0.59	0.24-1.43
Occupation						
Never employed in any risk occupation for lung cancer	138	163	1	–	1.00	–
Employed for over 1 year in a risk occupation	49	78	0.89	0.57-1.39	1.03	0.63-1.66
Cigarettes per day						
Never smokers	77	106	1	–	–	–
Moderate smokers (1-20)	74	100	1.11	0.70-1.75	–	–
Heavy smokers (> 20)	36	34	1.68	0.93-3.03	–	–
Duration of smoking						
Never smokers	77	105	1	–	–	–
1-30 years	44	68	1.22	0.72-2.06	–	–
Over 30 years	64	63	1.26	0.77-2.07	–	–
Total packets smoked over lifetime						
Never smokers	77	107	1	–	1	–
Moderate smokers	47	74	1.05	0.63-1.73	0.84	0.49-1.43
Heavy smokers	63	55	1.65	1.00-2.72	1.33	0.77-2.31
Alcohol consumption						
Non drinkers	29	79	1	–	1	–
Moderate drinkers (1-50 ml per day)	76	99	2.94	1.66-5.23	2.76	1.53-4.99
Heavy drinkers (> 50 ml per day)	80	80	5.16	2.80-9.49	4.83	2.55-9.14

OR: odds ratio; CI: confidence interval.

^aCrude model: adjusted for age and sex.

^bMultivariate model: adjusted for age, sex, education, occupation, total packets smoked over a lifetime and alcohol consumption.

for exclusion of individuals with previous respiratory diseases in the population-based study).

The most important limitation is the lower participation of population controls, a finding in line with most previous studies. Controls were not recruited simultaneously; instead, the population controls were enrolled from 1992 through 1995, and the hospital controls from 1999 through 2000. We feel, however, that the habits studied hardly varied in the interval between the two studies.

Conclusions

The ideal control group rarely exists in epidemiologic studies. In addition to theoretical work, empirical studies are needed to measure the impact of the different biases that can appear, so that these can be prevented and corrected. However, even though appropriate selection of controls is problematic, the most serious limitations can be avoided by bearing a series of basic

principles in mind⁸. Hospital controls are usually suitable where hospital cases are used, but could not represent the general population. Population controls, though generally preferable for reasons of validity, are expensive and difficult to obtain. Furthermore, for reasons of convenience, there is an important reason for not recruiting population controls, the need to ensure that the information bias that affects cases is similar⁹. In this study, while the characteristics of population and hospital controls were generally similar, differences were nevertheless observed in respect of alcohol consumption. Further studies are called for to ascertain whether there are differences between population or hospital controls, and for which diseases and populations it would be more appropriate to use one or the other.

Competing interests

The authors declare no conflict of interests.

Funding

This work has been partly funded by two investigation grants, one from the Fondo de Investigaciones Sanitarias (FIS) (Health Research Fund) (grant FIS 92/0176) and the Galician Regional Health Authority (grant XUGA 91010).

References

1. Rothman KJ. *Modern epidemiology*. Boston: Little Brown; 1986.
2. Wacholder S, McLaughlin JK, Silverman DT, et al. Selection of controls in case-control studies (II). Types of controls. *Am J Epidemiol*. 1992;135:1029-41.
3. Barros-Dios JM, Barreiro-Carracedo MA, Ruano-Ravina A, et al. Exposure to residential radon and lung cancer in Spain: a population-based case-control study. *Am J Epidemiol*. 2002;156:548-55.
4. Ruano-Ravina A, Figueiras A, Barros-Dios JM. Type of wine and risk of lung cancer. A case control study in Spain. *Thorax*. 2004;59:981-5.
5. Ahrens W, Merletti F. A standard tool for the analysis of occupational lung cancer in epidemiologic studies. *Int J Occup Environ Health*. 1998;4:236-40.
6. Miller AB. Hospital or population controls? It depends on the question. *Prev Med*. 1994;23:263-6.
7. Wacholder S, McLaughlin JK, Silverman DT, et al. Selection of controls in case-control studies (I). Principles. *Am J Epidemiol*. 1992;135:1019-28.
8. Wacholder S, McLaughlin JK, Silverman DT, et al. Selection of controls in case-control studies (III). Design options. *Am J Epidemiol*. 1992;135:1042-50.
9. Lasky T, Stolley PD. Selection of cases and controls. *Epidemiol Rev*. 1994;16:6-17.