

60. National Tobacco Information Online System. National Tobacco Information Online System (NATIONS). Country data [cited Accessed July 15 2002]. Available from: <http://apps.nccd.cdc.gov/nations>
61. Dixon MF, Genta RM, Yardley JH, Correa P. Classification and grading of gastritis. The updated Sydney System. International Workshop on the Histopathology of Gastritis, Houston 1994. *Am J Surg Pathol.* 1996;20:1161-81.
62. Dursun M, Yilmaz S, Yukselen V, Kilinc N, Canoruc F, Tuzcu A. Evaluation of optimal gastric mucosal biopsy site and number for identification of *Helicobacter pylori*, gastric atrophy and intestinal metaplasia. *Hepatogastroenterology.* 2004;51:1732-5.
63. Zhang C, Yamada N, Wu YL, Wen M, Matsuhisa T, Matsukura N. *Helicobacter pylori* infection, glandular atrophy and intestinal metaplasia in superficial gastritis, gastric erosion, erosive gastritis, gastric ulcer and early gastric cancer. *World J Gastroenterol.* 2005;11:791-6.
64. Eriksson NK, Farkkila MA, Voutilainen ME, Arkkila PE. The clinical value of taking routine biopsies from the incisura angularis during gastroscopy. *Endoscopy.* 2005;37:532-6.
65. Piantadosi S. Invited commentary: ecologic biases. *Am J Epidemiol.* 1994;139:761-4.
66. López AD, Collishaw NE, Piha T. A descriptive model of the cigarette epidemic in developed countries. *Tob Control.* 1994;3:242-7.
67. Leung WK, Lin SR, Ching JY, To KF, Ng EK, Chan FK, et al. Factors predicting progression of gastric intestinal metaplasia: results of a randomised trial on *Helicobacter pylori* eradication. *Gut.* 2004;53:1244-9.
68. Koizumi Y, Tsubono Y, Nakaya N, Kuriyama S, Shibuya D, Mat-suoka H, et al. Cigarette smoking and the risk of gastric cancer: a pooled analysis of two prospective studies in Japan. *Int J Cancer.* 2004;112:1049-55.
69. McFarlane GA, Wyatt J, Forman D, Lachlan GW. Trends over time in *Helicobacter pylori* gastritis in Kenya. *Eur J Gastroenterol Hepatol.* 2000;12:617-21.
70. Louw JA, Kidd MS, Kummer AF, Taylor K, Kotze U, Hanslo D. The relationship between *Helicobacter pylori* infection, the virulence genotypes of the infecting strain and gastric cancer in the African setting. *Helicobacter.* 2001;6:268-73.
71. Smith SI, Kirsch C, Oyedeji KS, Arigbabu AO, Coker AO, Bayerdoffer E, et al. Prevalence of *Helicobacter pylori vacA, cagA* and *iceA* genotypes in Nigerian patients with duodenal ulcer disease. *J Med Microbiol.* 2002;51:851-4.
72. Nessa J, Chart H, Owen RJ, Drasar B. Human serum antibody response to *Helicobacter pylori* whole cell antigen in an institutionalized Bangladeshi population. *J Appl Microbiol.* 2001;90:68-72.
73. El-Mahdi AM, Patchett SE, Char S, Domizio P, Fedail SS, Kumar PJ. Does *CagA* contribute to ulcer pathogenesis in a developing country, such as Sudan? *Eur J Gastroenterol Hepatol.* 1998;10:313-6.

---

## Comment. Gastric intestinal metaplasia and smoking

(Comentario. Metaplasia intestinal gástrica y tabaco)

Carlos A. González

Catalan Institute of Oncology, Barcelona, Spain.

Although the pathogenesis of intestinal metaplasia remains unknown<sup>1</sup>, it is well recognized that chronic gastritis and intestinal metaplasia are precursor conditions of gastric cancer. In intestinal metaplasia, the original gastric mucosa is replaced by glandular structures bearing an intestinal phenotype. Intestinal metaplasia is one of several steps in the prolonged precancerous process of gastric cancer, mostly of the intestinal histological type. There are 2 other types of intestinal metaplasia: the complete (type I) and the incomplete (type III) type. The incomplete type is more frequently associated with dysplasia and early carcinoma and is considered a more advanced stage of the process or of higher malignant potential<sup>2</sup>.

On the other hand, there is enough evidence about the causal relationship between smoking and gastric cancer risk<sup>3</sup>. The study by Peleteiro and colleagues deals with a systematic review of prevalence of intestinal metaplasia among *Helicobacter pylori* infected individuals and the correlation with na-

tional tobacco availability by means of an ecological analysis<sup>4</sup>.

Several case-control studies have shown the association between intestinal metaplasia and smoking. Although an ecological study is not the best appropriate epidemiological design to support the hypothesis of the role of smoking in the risk of intestinal metaplasia or in the progression from intestinal metaplasia to gastric cancer, there are no reviews published on the prevalence of intestinal metaplasia. This study provides a large, detailed, and useful information about the prevalence of intestinal metaplasia in 29 countries of five continents. Nevertheless, caution should be taken when the prevalence of intestinal metaplasia between studies and countries. The population source (clinical- or population-based), the age and the sex of participants, as well as the number of biopsies specimens, and the biopsy site, may influence the prevalence of intestinal metaplasia and therefore the comparability of results from different studies.

Information of tobacco availability in each country, expressed as the adult apparent tobacco consumption in cigarettes per day, were obtained from the Tobacco and Health Report of the WHO or from the National Tobacco Information Online System database, which provide reliable information at the population level. Despite the importance of these results, it is necessary to stress that causal relationships can be only derived from individual exposure data from studies to come.

## References

1. Akasak Y, Ishii T. Review: Histopathology and molecular pathology of intestinal metaplasia. *Curr Diag Pathol.* 2007;13:331-9.
2. Leung WK, Sung JJY. Review article: intestinal metaplasia and gastric carcinogenesis. *Aliment Pharmacol Ther.* 2002;16:1209-16.
3. Vineis P, Alavanja M, Buffler P, Fontham E, Franceschi S, Gao YT, et al. Tobacco and cancer: recent epidemiological evidence. *JNCI.* 2004;96:99-106.
4. Peleteiro B, Bastos J, Barros H, Lunet N. Systematic review of the prevalence of gastric intestinal metaplasia and its area-level association with smoking. *Gac Sanit.* 2008;22:236-46.