

Note from the field

Electronic clinical records in primary care for estimating disease burden and management. An example of COPD



Luis Verde-Remeseiro^a, Estrella López-Pardo^a, Alberto Ruano-Ravina^{b,c,*}, Francisco Gude-Sampedro^d, Ramón Castro-Calvo^e

^a Directive Staff, Santiago de Compostela University Clinic Hospital, Santiago de Compostela (A Coruña), Spain

^b Department of Preventive Medicine and Public Health, University of Santiago de Compostela (A Coruña), Spain

^c CIBER de Epidemiología y Salud Pública (CIBERESP), Spain

^d Epidemiology Unit, Santiago de Compostela University Clinic Hospital, Santiago de Compostela (A Coruña), Spain

^e Admission Unit, Santiago de Compostela University Clinic Hospital, Santiago de Compostela (A Coruña), Spain

ARTICLE INFO

Article history:

Received 4 February 2015

Accepted 23 March 2015

Available online 8 May 2015

Keywords:

Pulmonary disease

Chronic obstructive

Spirometry

Medical records systems

Computerized

Spain

ABSTRACT

Chronic obstructive pulmonary disease (COPD) is a significant health problem in developed countries. We aimed to estimate the prevalence of COPD in a single Spanish healthcare area. We also aimed to assess if there are any differences in prevalence and spirometry use among primary care services by utilizing already registered information. We designed a cross-sectional study to determine the prevalence of COPD and the performance of spirometries in each primary care service. A total of 8,444 patients were diagnosed with COPD, with a prevalence of 2.6% for individuals older than 39 years. The prevalence increased with age and was much higher in men. Significant heterogeneity was found in the prevalence of COPD and spirometry use among primary care services. COPD was underdiagnosed and there was wide variability in spirometry use in our area. Greater efforts are needed to diagnose COPD in order to improve its clinical outcomes and to refine registries so that they can be used as reliable sources of information

© 2015 SESPAS. Published by Elsevier España, S.L.U. All rights reserved.

La historia clínica electrónica en atención primaria en la estimación de la carga y la gestión de una enfermedad. El ejemplo de la EPOC

RESUMEN

La enfermedad pulmonar obstructiva crónica (EPOC) es un importante problema de salud en los países desarrollados. Se pretende estimar la prevalencia de la EPOC en un área sanitaria española para evaluar si existen diferencias en la prevalencia y en el uso de espirometrías entre los servicios de atención primaria utilizando información proveniente de registros. Se diseñó un estudio transversal para obtener la prevalencia de la EPOC y la realización de espirometrías en cada servicio de atención primaria. Hay 8444 pacientes diagnosticados de EPOC, con una prevalencia del 2,6% en mayores de 39 años. La prevalencia aumenta con la edad y es mucho más alta en los hombres que en las mujeres. Se observa una importante heterogeneidad en la prevalencia de la EPOC y en el uso de espirometrías entre servicios de primaria. La EPOC está infradiagnosticada y hay una fuerte variabilidad en el uso de espirometrías dentro de un área sanitaria. Deben hacerse más esfuerzos para diagnosticar la EPOC con el fin de mejorar los resultados clínicos, y refinar los registros para que puedan utilizarse como fuentes fiables de información sanitaria.

© 2015 SESPAS. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Enfermedad pulmonar obstructiva crónica

Espirometría

Historia clínica electrónica

España

Introduction

Chronic obstructive pulmonary disease (COPD) prevalence in Western countries is around 6–8%. In Spain, there is a prevalence of 10.2% in the population aged 40 to 80.¹ Prevalence would be even higher if patients older than 80 were taken into consideration.

Healthcare organizations have enforced strategies for improving the control and follow-up of COPD patients. Information and

communication technologies (ICTs) play an important role in its management. Using electronic health records a clinician can instantly have the patient's information and have access to his/her different tests. Furthermore, the disease prevalence can be calculated if the information is correctly registered.²

COPD prevalence information is scarce and there is a significant difference between registered prevalence and the true population prevalence. Many patients are not diagnosed because they do not have symptoms or clinicians are not aware of the value of spirometry. Spirometry is the core test for COPD, allowing the diagnosis of the disease and the assessment of its progression. At least one spirometry per year is recommended. Current guidelines

* Corresponding author.

E-mail address: alberto.ruano@usc.es (A. Ruano-Ravina).

Table 1
COPD prevalence broken down by age and gender in the Santiago de Compostela Health Area.

Variable	n	%
Total population	458,212	100
Females	236,630	51.6
COPD patients	8,444	1.84
≤38 years	181	0.10
39–64 years	1,829	1.10
65–74 years	2,025	4.43
≥75 years	4,409	7.79
Female COPD patients	3,012	1.27
≤38 years	77	0.08
39–64 years	584	0.7
65–74 years	593	2.44
≥75 years	1,758	5.01
Male COPD patients	5,432	2.45
≤38 years	104	0.11
39–64 years	1,245	1.50
65–74 years	1,432	6.69
≥75 years	2,651	12.31

recommend performing a spirometry on any smoker older than 40 who has smoked more than 10 pack-years.³ There is also a lack of information on how resources are used in primary care facilities to manage the disease. Some investigations have observed that a significant percentage of COPD patients have not had a spirometry performed in the last year and some COPD patients have even been diagnosed without this test.⁴

We aim to estimate COPD prevalence and the rate of spirometries performed in primary health care services using data exclusively obtained from health information systems.

Subjects and methods

The study was performed in the Santiago de Compostela (A Coruña, Spain) health area. It has a population of 458,212 inhabitants, with 22.6% older than 64. Primary care is provided by 21 health services comprising 72 primary care facilities.

We designed a cross-sectional study to obtain COPD prevalence and the performance of spirometries during 2012 in the public health system. It covers approximately 97% of the entire Galician population. Data extraction was performed during June 2013. To retrieve COPD patients we used the code R95 of the International Classification of Primary Care-2. Health managers have access to all information systems, though it is necessary to have skills and training to retrieve specific information from the databases. We have also obtained COPD prevalence and performance of spirometries per each primary care service. We calculated non-parametric correlations (Spearman's rho) to assess the relationship between COPD prevalence and the number of spirometries per 100 COPD patients for each primary care service with SPSS v17.

Results

In **Table 1** there is a sample description. 8,444 patients had COPD and prevalence was 1.8% (2.6% for those older than 39). COPD prevalence in individuals older than 74 was nearly twofold that of the 65–74 age group. COPD prevalence in males was double that of females. COPD prevalence for males 75 or older was 12.3% and 5.0% for females.

In 2012, a total of 7,071 spirometries were performed on COPD patients in the healthcare area. 14.9% of COPD patients had a spirometry registered in primary or specialized care. 71.2% of all spirometries were performed at a hospital and the remaining in family practices (n = 1,986).

Table 2
COPD prevalence for primary care services and spirometries performed per 100 COPD patients during 2012.

Primary care service	COPD prevalence in individuals older than 39	Spirometries per 100 COPD patients older than 39
1	1.74	0.53
2	1.88	22.54
3	1.88	27.70
4	1.96	2.19
5	1.98	35.11
6	2.17	5.69
7	2.24	6.60
8	2.26	----
9	2.31	35.34
10	2.40	30.33
11	2.44	19.83
12	2.48	10.79
13	2.56	33.57
14	2.60	9.40
15	3.22	42.69
16	3.24	----
17	3.34	22.22
18	3.53	28.45
19	3.77	30.56
20	4.09	12.69
21	4.75	----
Total	2.62	20.28

Table 2 shows the prevalence of COPD per each primary care service with the corresponding number of spirometries performed per 100 COPD patients. The primary service with the highest prevalence is three times that of the service with the lowest prevalence (1.74% vs. 4.75%), with a similar age distribution. There was no correlation between COPD prevalence and the number of spirometries performed in each primary care service (Spearman's $\rho = 0.31$; $p = 0.21$). Three primary care services did not perform any spirometries in 2012. 27% of COPD patients had at least one hospital admission during 2012.

Discussion

To our knowledge, this is the first study performed in Spain using only registry-based information to calculate COPD prevalence. It is also the first to compare if there are differences in the prevalence and the ratio of spirometries performed per patient among primary care services. Using health information systems we have observed that COPD is clearly underdiagnosed in our area and also that spirometry is clearly underused in COPD patients. Furthermore, there is a great heterogeneity in its use among primary care facilities, but this heterogeneity does not depend on COPD prevalence. Low COPD prevalence in our area denotes underdiagnosis, in line with other studies. The EPI-SCAN study found 70% of unknown disease in the general population.¹ Other studies have shown 3.5% COPD prevalence based on clinical registries⁵ while ours finds a 2.6% prevalence, which is lower than expected.

The exhaustiveness and reliability of disease registration in clinical databases is a critical aspect when using health information systems to obtain epidemiological information. If this information is appropriately registered it could provide important information for health managers and it would be unnecessary to carry out studies specifically directed to estimate the prevalence of many diseases. Clinicians should be aware of the importance of this registration and health administration should facilitate it.

We have found significant differences between primary care facilities. In some cases, despite the availability of spirometers, they are not used in family practices. It has not been until recently that spirometry results have been included in electronic health records.

In the area of study all hospital-based spirometries are registered but this might not be the case for spirometries performed in family practices. If spirometries are performed during a medical visit, they can be registered or not, and therefore spirometries performed in primary care might be underestimated in the present study. It is quite common that COPD patients do not have spirometric values in their clinical history. The AUDIPOC study, a clinical audit including more than 5,000 COPD patients, followed up for three months to assess hospital readmissions and mortality, and observed that there were significant differences in the availability of spirometric data.⁶ Spirometry was registered only in 41%,⁷ 59%⁸ and 30%⁹ of patients diagnosed with COPD in other investigations.

The main advantage of this study is that we have not used a sample but rather the entire population of the Santiago de Compostela health area, and therefore a possible selection bias should not be present. The main shortcoming is that this research is based on registered information and registration strongly depends on the physician's interest or responsibility. This is reflected by the high variability in COPD prevalence considering different primary care services.

COPD prevalence is underdiagnosed and the number of spirometries performed in primary care must be significantly increased in our area. It is very important to diagnose and treat occult COPD. The present study also demonstrates that health information systems can be used for multiple purposes, from knowing the prevalence of many diseases to having information regarding disease management. These registries might also be useful in the surveillance and monitoring of different health problems.¹⁰ The usefulness of this information resides in the physicians registration of medical data and health administration should allocate resources to strengthen registration.

What is known about the topic?

Chronic obstructive pulmonary disease (COPD) is one of the most significant chronic diseases. Spirometry is the core test for COPD. Health information systems could be used to estimate the frequencies of many diseases and in the present paper we use them to estimate COPD prevalence and spirometry use within a healthcare area.

What does this study add to the literature?

COPD is underdiagnosed and there are significant differences in spirometry use within one healthcare area. Integrated health information systems can be successfully used to compare the performance of primary care practices and to estimate the frequency of chronic diseases.

Editor in charge

Pere Godoy.

Transparency declaration

The corresponding author on behalf of the other authors guarantee the accuracy, transparency and honesty of the data and information contained in the study, that no relevant information has been omitted and that all discrepancies between authors have been adequately resolved and described.

Authors' contributions

A. Ruano-Ravina, L. Verde-Remeseiro and E. López-Pardo designed the research and interpreted the results. R. Castro-Calvo and F. Gude-Sampedro extracted the information and merged the databases. All authors wrote the paper and analyzed the results. All authors gave important insights to the work and approved the final version of the manuscript. A. Ruano-Ravina is the guarantor of the paper.

Funding

This research has been sponsored by an unrestricted grant by Almirall. The sponsor did not have any role in the design, writing or interpretation of the present research.

Conflicts of interest

None of the authors are employees of Almirall or have a financial relationship with this Company. One of the authors (A. Ruano-Ravina) belongs to the GACETA SANITARIA editorial committee, but was not involved in the editorial process of the manuscript.

References

1. Soriano JB, Miravittles M, Borderías L, et al. Geographical variations in the prevalence of COPD in Spain: relationship to smoking, death rates and other determining factors. *Arch Bronconeumol.* 2010;46:522–30.
2. Chao A, Tsay PK, Lin SH, et al. The applications of capture-recapture models to epidemiological data. *Stat Med.* 2001;20:3123–57.
3. Ancochea Bermúdez J, editor. Estrategia en EPOC del Sistema Nacional de Salud [Internet]. Madrid: Ministerio de Sanidad; 2009. [Cited 2015 Feb 26]. 183 p. Available from: <http://www.msssi.gob.es/organizacion/sns/planCalidadSNS/docs/EstrategiaEPOCSNS.pdf> página 17.
4. Pellicer Císcar C, Soler Cataluña JJ, Andreu Rodríguez AL, et al. On behalf of Grupo EPOC de Sociedad Valenciana de Neumología. Diagnosis of COPD in hospitalised patients. *Arch Bronconeumol.* 2010;46:64–9.
5. Ställberg B, Janson C, Johansson G, et al. Management, morbidity and mortality of COPD during an 11-year period: an observational retrospective epidemiological register study in Sweden (PATHOS). *Prim Care Respir J.* 2014;23:38–45.
6. Pozo-Rodríguez F, López-Campos JL, Álvarez-Martínez CJ, et al. Clinical audit of COPD patients requiring hospital admission in Spain: AUDIPOC study. *PLoS One.* 2012;7:e42156.
7. Maio S, Baldacci S, Martini F, et al. COPD management according to old and new GOLD guidelines: an observational study with Italian general practitioners. *Curr Med Res Opin.* 2014;30:1033–42.
8. Arne M, Lisspers K, Ställberg B, et al. How often is diagnosis of COPD confirmed with spirometry. *Respir Med.* 2010;104:550–6.
9. Caramori G, Bettoncelli G, Carone M, et al. Degree of control of physician-diagnosed asthma and COPD in Italy. *Monaldi Arch Chest Dis.* 2007;67:15–22.
10. Esteban-Vasallo MD, Domínguez-Berjón MF, Astray-Mochales J, et al. Epidemiological usefulness of population-based electronic clinical records in primary care: estimation of the prevalence of chronic diseases. *Fam Pract.* 2009;26:445–54.