

## Brief original article

## Factors associated with early childhood caries in Chile



Lorena Hoffmeister, Patricia Moya\*, Carolina Vidal, Dafna Benadof

Escuela de Salud Pública, Universidad Mayor, Santiago, Chile

## ARTICLE INFO

## Article history:

Received 7 July 2015

Accepted 30 September 2015

Available online 3 December 2015

## Keywords:

Dental caries

Child

Preschool

Socioeconomic factors

## ABSTRACT

**Objective:** To determine the prevalence of caries and identify the associated factors among children aged 2 and 4 years in southern Chile.

**Methods:** A cross-sectional study was conducted in a random sample of 2,987 children. Dental examinations were performed in all participating children and a self-administered survey was administered to their primary caregivers. The statistical analysis included a zero inflated negative binomial regression model.

**Results:** The prevalence of caries was 20.3% for 2-year-olds and 52.7% for 4-year-olds. The factors associated with caries included the following: living in the Bio-Bío region,  $PR_{2\text{year-old}}: 1.65 (1.10-2.47)$ ,  $PR_{4\text{year-old}}: 1.44 (1.18-1.75)$  and having a low socioeconomic position,  $PR_{2\text{year-old}}: 5.39 (2.14-13.57)$ ,  $PR_{4\text{year-old}}: 3.71 (2.68-5.13)$  or a medium socioeconomic position,  $PR_{2\text{year-old}}: 2.79 (1.06-7.33)$ ,  $PR_{4\text{year-old}}: 2.70 (1.92-3.79)$ . In 4-year-olds, a high frequency of consuming sugary drinks at bedtime ( $PR: 1.30; 1.06-1.59$ ), not brushing teeth daily ( $PR: 1.37; 1.18-1.60$ ), and the presence of gingivitis ( $PR: 1.64; 1.28-2.09$ ) were also associated with caries.

**Conclusions:** The high prevalence of caries in this young group is concerning. The factors associated with caries identified in this study could help in the design of preventive interventions at early ages.

© 2015 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Factores asociados con la caries infantil precoz en Chile

## RESUMEN

**Objetivo:** Determinar la prevalencia y los factores asociados a la caries en niños y niñas de 2 y 4 años de edad en el sur de Chile.

**Métodos:** Estudio transversal, con muestra probabilística de 2987 niños y niñas. Se realizaron exámenes dentales a los participantes y una encuesta autoadministrada a sus cuidadores. El análisis estadístico incluyó un modelo binomial negativo cero inflado.

**Resultados:** La prevalencia de caries fue del 20,3% (2 años de edad) y del 52,7% (4 años de edad). Los factores asociados a la caries fueron vivir en la región del Bio-Bío,  $RP_{2\text{años}}: 1,65 (1,10-2,47)$ ,  $RP_{4\text{años}}: 1,44 (1,18-1,75)$ , y la posición socioeconómica baja,  $RP_{2\text{años}}: 5,39 (2,14-13,57)$ ,  $RP_{4\text{años}}: 3,71 (2,68-5,13)$ , o media,  $RP_{2\text{años}}: 2,79 (1,06-7,33)$ ,  $RP_{4\text{años}}: 2,70 (1,92-3,79)$ . En los niños y las niñas de 4 años de edad, la alta frecuencia de consumo de bebestibles azucarados antes de dormir ( $RP: 1,30; 1,06-1,59$ ), no lavarse los dientes a diario ( $RP: 1,37; 1,18-1,60$ ) y la gingivitis ( $RP: 1,64; 1,28-2,09$ ) se asociaron a tener caries.

**Conclusiones:** Esta alta prevalencia de caries es preocupante. Los factores asociados a la caries identificados pueden ayudar en el diseño de intervenciones preventivas en etapas precoces.

© 2015 SESPAS. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Palabras clave:

Caries dental

Preescolar

Factores socioeconómicos

## Introduction

Early childhood caries is a significant public health problem due to its high prevalence in economically disadvantaged groups.<sup>1</sup> Factors such as consumption of sugary foods, dental plaque accumulation, and number of people in the household have been found to be associated with caries.<sup>2,3</sup>

Few studies have evaluated oral health and socioeconomic disparities in preschool children in Latin American countries.<sup>4,5</sup> In Chile, efforts have been made to diminish health disparities (e.g.,

water fluoridation programs). Nevertheless, the caries distribution associated with socioeconomic, dietary and oral hygiene factors has not been described for preschool children.

We aimed to identify factors associated with caries extent and estimate the prevalence of caries among preschool children in southern Chile.

## Methods

## Design and sampling process

The following was an observational, cross-sectional study. The sampling frame was the daycare centers register ( $N=2,473$ ). A stratified multistage probability sampling method was used.

\* Corresponding author.

E-mail address: [salud publica@umayor.cl](mailto:salud publica@umayor.cl) (P. Moya).

Stratification was based on age (2 and 4-year-old) and geographical region (Bio-Bío, Araucanía, Los Lagos-Los Ríos, and Aysén-Magallanes). The sampling unit was daycare centers in each stratum. We included children attending all three types of daycare centers (public without out-of-pocket charges, subsidized, and private).

This study was approved by the Bioethics Committee of Universidad Mayor, Chile.

#### *Study variables and survey*

Clinical variables included the traditional dental caries index (dmft),<sup>6</sup> caries prevalence (dmft  $\geq 1$ ), and gingivitis (only measured in 4-years-olds). A socioeconomic position index was created using educational level of the head of the household and household income variables. Sociodemographic variables included gender, age, educational level, and region of residence; family composition was assessed with the number of children and adults living in the household.<sup>2,7</sup> Nutritional factors were evaluated with bedtime sugary beverages frequency, tea consumption (because of its fluoride composition), and persistence of night-time breastfeeding. Oral hygiene habits included tooth brushing frequency and adult supervision. A self-administered questionnaire for caregivers evaluated these variables.

#### *Data collection*

Data was collected between October 2009 and May 2010. Caregivers of children aged 2 and 4 were sent a letter describing the study and asking for consent to examine their child. The letter also included a survey for the caregivers to fill out at home. Children of parents who consented underwent a standardized dental examination<sup>6</sup> performed by two trained dentists. Calibration was conducted in 20 children. The caries index kappa coefficient for inter-rater agreement was  $K = 0.89$  and  $K_1 = 0.98$ ,  $K_2 = 0.91$  for intra-rater agreement. The gingivitis kappa coefficient was  $K = 0.773$  for inter-rater agreement and  $K_1 = 1.00$ ,  $K_2 = 1.00$  for intra-rater agreement.

#### *Data analysis*

According to the sample design and to obtain representative results by region, sampling weights were calculated as the inverse probability of being included in the study. Frequency tables, point estimates and 95% confidence intervals were used. The prevalence of dental caries, dmft index, and Significant Caries Index (SiC, which identifies the third most affected by caries in the population) were calculated for each stratum.<sup>8</sup>

We used a zero inflated regression model to analyze the association of sociodemographic factors, family composition, nutritional and oral habits with the dmft index for 2 and 4-year-old children. The association of gingivitis with the dmft index was evaluated only in 4-year-olds. We used the Vuong test to select between the Poisson (ZIP) and negative binomial (ZINB) distributions. Data analysis was performed using Stata 13.

### **Results**

The final sample size was 1,387 two-year-old children and 1,600 four-year-old children. The response rate was 75.4% and 76.2% respectively. A total of 65.7% of the 2-year-olds and 59.1% of the 4-year-olds were from families with low socioeconomic position. The socioeconomic position distribution was similar between regions (data not showed). The highest educational level reported by 65.4% of the 2-year-olds' caretakers and 63.8% of those of 4-year-olds was secondary school or less.

Caries prevalence in 2-years-olds was 20.3% and 52.7% in 4-year-olds. In both groups, the prevalence of caries was lower as the household's socioeconomic position category was higher (Table 1).

The mean dmft was 0.6 in 2-year-olds (95% confidence interval [95%CI]: 0.5-0.8). In this group, the mean number of decayed teeth was 0.6 (95%CI: 0.5-0.7) and mean number of filled teeth was 0.0 (95%CI: 0.0-0.1). No extractions were found. The mean dmft was higher in children from low socioeconomic position households than in those from high socioeconomic position households (Table 1). No statistically significant differences between boys and girls were found for the dmft index (p-value for U Mann-Whitney test was 0.097).

In 4-year-olds, the mean dmft was 2.5. This group had a mean number of decayed teeth of 2.0 (95%CI: 1.7-2.4), mean number of filled teeth of 0.4 (95%CI: 0.3-0.6), and mean number of extracted teeth of 0.0 (95%CI: 0.0-0.1). The mean dmft was 3.2 in 4-year-olds from low socioeconomic position households, compared with 0.6 in those from high socioeconomic position households.

The SiC index was 1.8 for 2-years-old children and 6.0 for 4-years-olds. In both age groups, children from low socioeconomic position households had higher SiC indexes than children from other socioeconomic position households.

Table 2 shows the results obtained from the multivariable model for dmft. For 2-year-old children living in the Bio-Bío region, the dmft was 1.65 times larger than it was for those living in the region of Aysén-Magallanes, whereas for 4-years-old it was 1.44 times as large. Children in low and middle socioeconomic position households had a significantly larger dmft index than those in higher socioeconomic position households.

Four-year-old children who drank sugary beverages before bedtime with high frequency had a dmft 1.30 times as large as those who drank before bedtime less than 3 times per week. Four-year-olds who did not brush their teeth daily had a dmft 1.37 times larger than those who brushed their teeth daily. In this age group gingivitis was associated with the dmft index (PR = 1.64).

### **Discussion**

Our findings reveal a high prevalence of dental caries in the preschool children in Chile. Countries such as the United States<sup>9</sup> and Brazil<sup>4</sup> have reported lower caries prevalences among comparable populations. Despite these differences, caries showed a similar distribution pattern among socioeconomic levels as described in other studies.<sup>4,9</sup> It is important to note that the dental caries index used for this study followed the guidelines of the WHO,<sup>6</sup> which does not differentiate between stages of development of dental cavities. This situation might underestimate the caries prevalence in this population.

The mean dmft in 4-year-olds was lower than that reported in similar studies in Chile,<sup>10</sup> but higher than that reported in similar populations in other Latin American countries<sup>11</sup> and indicates considerable oral damage due to caries. Health disparities can be further understood with the SiC index which showed that children in low socioeconomic position households had extensive caries experience.

The multivariable and descriptive analyses confirmed large-scale inequalities in socioeconomic position in both age groups, which is consistent with the literature describing older children.<sup>1,12</sup> These inequalities are reflected in the descriptive results, which showed that 2 year old children of low socioeconomic groups had 10 times more caries than those of higher socioeconomic groups. Also the multivariable model showed a prevalence ratio of caries in children of low socioeconomic groups of at least twice that of higher socioeconomic groups. On the other hand, nutritional factors

**Table 1**

Summary statistics for the caries prevalence (%), dental caries index mean (dmft), and significant caries index mean (SiC) according to socioeconomic position categories (SEP), in 2 and 4 year-olds, southern Chile.

	2 years		4 years	
	n	Value (95%CI)	n	Value (95%CI)
<i>Prevalence of caries (%)</i>				
Low SEP	198/902	24.5 (21.0-28.4)	544/944	58.8 (51.0-66.2)
Middle SEP	34/280	11.4 (7.6-16.8)	172/374	51.6 (41.1-62.0)
High SEP	4/102	2.3 (0.6-7.9)	37/176	27.3 (11.8-51.4)
All groups	236/1,284	20.3 (17.3-23.2)	753/1,494	52.7 (46.8-58.5)
<i>dmft (mean)</i>				
Low SEP	902	0.8 (0.7-1.0)	944	3.2 (2.5-3.8)
Middle SEP	280	0.3 (0.1-0.4)	374	1.8 (1.4-2.3)
High SEP	102	0.1 (0.0-0.2)	176	0.6 (0.3-0.9)
All groups	1,284	0.6 (0.5- 0.8)	1,494	2.5 (2.1-3.0)
Low SEP	271	2.1 (1.7-2.4)	375	7.1 (6.4-7.9)
<i>SiC (mean)</i>				
Middle SEP	56	0.7 (0.4-1.0)	105	4.6 (4.0-5.2)
High SEP	28	0.3 (0.0-0.6)	19	1.2 (0.5-1.9)
All groups	355	1.8 (1.5-2.0)	499	6.0 (5.3-6.7)

95%CI: 95% confidence interval.

<sup>a</sup> Weighted analyses.

contributed independently to caries prevalence. We hypothesize that these behaviors replaced previous behaviors also associated with nighttime feeding at earlier ages. Oral hygiene habits were found to be protective factors at age 4 which might be due in part to increased motor control,<sup>13</sup> and their increased ability to understand instructions.<sup>14</sup> Other factors such as parents educational level and dental education in daycare centers might also affect oral hygiene among individuals, but they could not explain

**Table 2**

Zero inflated negative binomial regression models for dental caries index (number of teeth with history of caries) in children aged 2 and 4 years old.

	2 years PR (95%CI)	4 years PR (95%CI)
<i>Socio-demographics</i>		
Region: Aysén-Magallanes	1.00	1.00
Region: Bio-Bío	1.65 (1.10-2.47)	1.44 (1.18-1.75)
Region: Araucanía	1.27 (0.823-1.96)	0.88 (0.72-1.07)
Region: Los Ríos- Los Lagos	0.65 (0.40-1.03)	0.94 (0.75- 1.17)
Household SEP: high	1.00	1.00
Household SEP: low	5.39 (2.14-13.57)	3.71 (2.68-5.13)
Household SEP: middle	2.79 (1.06-7.33)	2.70 (1.92-3.79)
Young maternal age (<21 years)	1.32 (0.87-2.02)	1.11 (0.73-1.67)
≥2 children living in the household	0.89 (0.65-1.21)	1.02 (0.87-1.20)
≥3 adults in household	0.99 (0.74-1.33)	1.06 (0.92-1.23)
Gender of child (girl)	0.96 (0.71-1.29)	0.89 (0.77-1.02)
<i>Nutritional habits</i>		
High frequency of sugary drinks at bedtime <sup>a</sup>	1.43 (0.97-2.10)	1.30 (1.06-1.59)
High frequency of juices or fizzy drinks <sup>a</sup>	1.14 (0.79-1.63)	1.08 (0.88-1.32)
High frequency of nocturnal bottle feeding <sup>a</sup>	0.95 (0.70-1.30)	0.97 (0.84-1.13)
High frequency of breast feeding <sup>a</sup>	1.33 (0.93-1.91)	1.07 (0.74-1.52)
High frequency of tea drinking <sup>a</sup>	0.81 (0.59-1.10)	1.07 (0.92-1.23)
<i>Oral hygiene</i>		
Not brushing teeth daily <sup>b</sup>	0.96 (0.71-1.29)	1.37 (1.18-1.60)
Supervision of brushing by an adult <sup>c</sup>	1.09 (0.81-1.48)	0.95 (0.82-1.10)
Gingivitis (yes) <sup>d</sup>	-	1.64 (1.28-2.09)

95%CI: 95% confidence interval; SEP: socioeconomic position.

Reference categories:

<sup>a</sup> Less than three times a week.

<sup>b</sup> Brushing teeth daily.

<sup>c</sup> No supervision by adult.

<sup>d</sup> Absence of gingivitis.

the difference between ages. The higher risk of developing caries in the region of Bio-Bío could be explained by the extremely low levels of fluoridated water in this area (0.003 to 0.055 ppm).<sup>15</sup>

Our results showed that there were no statistically significant differences between boys and girls in the number of carious lesions. The literature regarding this topic for this age group is scarce and not conclusive.<sup>16</sup>

The study limitations are that all sampled children attended daycare facilities, affecting the results' external validity; also the cross-sectional study design emphasizes the need to be cautious with the interpretation of its results. It is important to consider that nutritional and oral hygiene variables might be influenced by oral health status and therefore we cannot establish these as risk factors for dental caries.

On the other hand, the strength of this study relies on the sample size and on the heterogeneity of the sample in terms of geographical distribution, family characteristics, and type of daycare centers (public without out-of-pocket charges, subsidized, and private).

In conclusion, this study found a high caries prevalence in children at very young ages in this region of Latin America. Socioeconomic position was found to be associated with caries experience. Nutritional habits were associated with caries development in all children, independent of their socioeconomic status and region of residency. We suggest that other factors such as family traditions might be related to dietary habits in this population. We encourage the implementation of oral health education and prevention programs aimed at improving nutritional habits at a familial level so that children can experience a healthy home environment.

## Editor in charge

Glòria Pérez.

## Transparency declaration

The corresponding author on behalf of the other authors guarantee the accuracy, transparency and honesty of the data and information contained in the study, that no relevant information has been omitted and that all discrepancies between authors have been adequately resolved and described.

**What is known about the topic?**

Early childhood caries is a significant public health problem that greatly affects children's quality of life. In the Americas the impact of income inequalities affects people's oral health.

**What does this study add to the literature?**

This study provides insight into the oral health of children aged 2 and 4 years in Chile. This study is among the first in Latin America to evaluate the relationship between caries and socioeconomic, dietary and oral hygiene in this group.

**Authorship contributions**

L. Hoffmeister and P. Moya were involved in the conception and design of the work. C. Vidal carried out the analysis and interpretation of data. D. Benadof contributed to the writing and its critical review with important intellectual contributions. All authors have approved the final version and are jointly accountable for all aspects of the work.

**Funding**

The Ministry of Health of Chile contributed to the accomplishment of this work.

**Conflicts of interest**

None.

**References**

1. Antunes JLF, Narvai PC, Nugent ZJ. Measuring inequalities in the distribution of dental caries. *Community Dent Oral Epidemiol.* 2004;**32**:41–8.
2. Harris R, Nicoll AD, Adair PM, et al. Risk factors for dental caries in young children: a systematic review of the literature. *Community Dent Health.* 2004;**21**(Suppl 1):71–85.
3. Moimaz SAS, Fadel CB, Lolli LF, et al. Social aspects of dental caries in the context of mother-child pairs. *J Appl Oral Sci.* 2014;**22**:73–8.
4. Dos Santos Junior VE, de Sousa RMB, Oliveira MC, et al. Early childhood caries and its relationship with perinatal, socioeconomic and nutritional risks: a cross-sectional study. *BMC Oral Health.* 2014;**14**:47.
5. Costa LR, Daher A, Queiroz MG. Early childhood caries and body mass index in young children from low income families. *Int J Environ Res Public Health.* 2013;**10**:867–78.
6. World Health Organization. *Oral Health Surveys: Basic Methods.* 4th ed. Geneva: WHO; 1997. 67 p.
7. Antunes JLF, Frazao P, Narvai PC, et al. Spatial analysis to identify differentials in dental needs by area-based measures. *Community Dent Oral Epidemiol.* 2002;**30**:133–42.
8. Bratthall D. Introducing the Significant Caries Index together with a proposal for a new global oral health goal for 12-year-olds. *Int Dent J.* 2000;**50**:378–84.
9. Dye B, Tan S, Smith V, et al. Trends in oral health status: United States, 1988–1994 and 1999–2004. *Vital Health Stat.* 2007;**11**:1–92.
10. Zaror Sánchez C, Pineda Toledo P, Orellana Cáceres JJ. Prevalencia de caries temprana de la infancia y sus factores asociados en niños chilenos de 2 y 4 años. *Int J Odontostomatol.* 2015;**5**:171–7.
11. Carvalho F, Carvalho CRB, Xavier A, et al. Dental caries experience in preschool children of Bauru, SP, Brazil. *Brazilian Journal of Oral Science.* 2009;**8**.
12. Delgado-Angulo EK, Hobdell MH, Bernabé E. Poverty, social exclusion and dental caries of 12-year-old children: a cross-sectional study in Lima, Peru. *BMC Oral Health.* 2009;**9**:16.
13. Gerber RJ, Wilks T, Erdie-Lalena C. Developmental milestones: motor development. *Pediatr Rev.* 2010;**31**:267–76.
14. Wilks T, Gerber RJ, Erdie-Lalena C. Developmental milestones: cognitive development. *Pediatr Rev.* 2010;**31**:364–7.
15. Gomez S, Fernandez O. [Water fluoridation experience in Chile]. In: Gomez S, editor. [Fluoride therapy in dentistry]. 4th ed. 2010. p. 211–21.
16. Kramer AA, Hakeberg M, Petzold M, et al. Demographic factors and dental health of Swedish children and adolescents. *Acta Odontol Scand.* 2015;**1**:1–8.