

Review

Salutogenic interventions and health effects: a scoping review of the literature



Óscar Suárez Álvarez^{a,b,*}, María Teresa Ruiz-Cantero^{c,d}, Viola Cassetti^e, Rafael Cofiño^b, Carlos Álvarez-Dardet^{c,d}

^a Doctoral Programme in Health Science, University of Alicante, Alicante, Spain

^b Public Health Directorate, Asturias Regional Ministry of Health, Oviedo, Spain

^c Research Group in Public Health, University of Alicante, Alicante, Spain

^d CIBER de Epidemiología y Salud Pública (CIBERESP), Spain

^e School for Health and Related Research (SCHARR), University of Sheffield, Sheffield, United Kingdom

A B S T R A C T

Keywords:

Salutogenesis
Public health
Systematic review
Health assets
Health outcomes

Objective: To explore the typology of implemented salutogenic interventions and the health effects described by the authors.

Method: A scoping review of the literature published (PubMed, Embase, Web of Science and Scopus) over the last ten years (2007-2016) was conducted. Articles that included interventions with assessments of health outcomes were selected for this review. The kappa index (86.4%) was calculated for the classification and extraction of information.

Results: 61 papers were selected out of the 676 works identified. These were categorized into individual, grouped, mixed and intersectoral interventions. A total of 85% of the interventions described positive effects. Adverse effects were not reported. Methodological limitations were identified in 75% of the papers. The interventions addressed a large variety of topics, especially in the field of mental health and chronic diseases. Collective actions described more positive effects. Intersectoral actions were the only type of interventions to have approached the impact of mortality reduction.

Conclusions: Findings support the existence of positive health effects of salutogenic approaches. Important methodological limitations were identified, such as biases in the selection of participants and sample sizes. Studies need to be conducted with improved monitoring and evaluation designs. A more robust theoretical framework and tools to evaluate the salutogenic contents are needed.

© 2020 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Las intervenciones salutogénicas y los efectos en la salud. Una scoping review de la literatura científica

R E S U M E N

Palabras clave:

Salutogénesis
Salud pública
Revisión sistemática
Activos de salud
Resultados de salud

Objetivo: Conocer el tipo de las intervenciones salutogénicas identificadas en la literatura científica y los efectos descritos por los autores en la salud de la población.

Método: Se realizó una scoping review de trabajos publicados entre los años 2007 y 2016 en español, inglés y francés en PubMed, Embase, Web of Science y Scopus. Se seleccionaron y clasificaron trabajos de investigación que contemplaran una intervención con evaluación de resultados en salud. Se calculó el índice kappa (86,4%) para la clasificación y la extracción de la información.

Resultados: Se seleccionaron 61 trabajos de investigación entre 676 trabajos identificados. Se clasificaron en intervenciones individuales, grupales, mixtas e intersectoriales. El 85% de las intervenciones describieron efectos positivos. No se encontraron efectos adversos. Se observaron limitaciones metodológicas en el 75% de los trabajos. Las intervenciones abordaban gran variedad de temas, especialmente en el campo de la salud mental y las enfermedades crónicas. Las intervenciones colectivas describen más efectos positivos. Las actuaciones intersectoriales representan el único tipo de intervenciones que ha abordado el impacto en la reducción de la mortalidad.

Conclusiones: Los hallazgos descritos sugieren la existencia de efectos positivos para la salud. Se identifican importantes limitaciones metodológicas. Es necesario desarrollar estudios en mejores condiciones de control, así como marcos teóricos más robustos y herramientas que permitan evaluar el contenido salutogénico de las intervenciones.

© 2020 SESPAS. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: Oscarsuarezalvarez@yahoo.es (Ó. Suárez Álvarez).

Introduction

In 1971, the medical sociologist Aaron Antonovsky presented the findings of a study¹ he conducted on a group of women who had survived Nazi concentration camps. He found that some of these women, despite having suffered stressful situations, had maintained good physical and mental health. Antonovsky was interested in the factors that helped these women to sustain good health, an approach which led him to develop what is today known as salutogenic theory.^{2,3}

One of his major contributions was to introduce this theoretical framework as a new paradigm of positive health, in contrast to the pathogenic approach. This theoretical framework supports practices aimed at promoting the enhancement and better use of individual and collective resources to improve population health and well-being.⁴

Antonovsky was one of the first authors to develop an approach which allows studying and quantifying positive protective health factors and their impact on the population. His theories, despite having been applied to more individual approaches with the development of concepts such as sense of coherence (SOC) or generalized resistance resources (GRRs), have also influenced other fields aimed at population-wide approaches.⁵ Antonovsky introduced the salutogenic concept of SOC as a specific way of viewing life as comprehensible, manageable and meaningful.⁶ On the other hand, GRRs could be considered as those elements related to the material, biological, psychological and social means that would make it easier for people to perceive life as comprehensible, manageable and meaningful. GRRs would thus act as a kind of health asset that could help improving or maintaining the health of the individuals and the communities in which they live. They act on the SOC and constitute one of the main elements of our study.

Based on this theoretical proposal to carry out the development of possible empirical applications, some authors⁷ propose that salutogenic interventions should be designed to enable communities to create shared life visions and to be part of decision making (meaningfulness); develop shared mental models about the change process and desired outcomes (comprehensibility); enable communities to identify life demands (e.g., stressors, challenges) and GRRs that need to be balanced (manageability) as well as life opportunities (e.g., assets, learning situations) that stimulate health development.

Other authors have developed theoretical models to implement the salutogenic paradigm in different sectors, such as schools,⁸ communities or neighbourhoods,⁵ health systems,⁹ or within different population groups such as children¹⁰ or persons with chronic diseases.¹¹ Some interventions⁸ have proposed new ways of studying population health by incorporating a positive view of health, an assets orientation, emphasizing both the individual level –by measuring the SOC– and the social or community levels by researching concepts like social capital.

In 2006, Eriksson and Lindström¹² carried out a systematic review of 471 papers to study the relation between perceived health and SOC. They found that the stronger the SOC, the better the perceived health. A study recently conducted in Spain¹³ based on tracking periods over 8, 12 and 20 years, revealed links between perceived health and mortality. The SOC was found to be indirectly associated to a lower mortality rate. In a more social dimension, Holt-Lunstad¹⁴ performed a meta-analysis of 148 papers and observed that people with stronger social networks had a 50% increased likelihood of survival, even after adjusting for risk factors. This means that individuals with a stronger SOC or living in communities with greater social capital have better health outcomes.

Despite the numerous empirical applications of the salutogenic theory in the field of public health, reviews of salutogenic-based interventions and their effects on population health are limited.

This has resulted in a range of fragmented actions under the salutogenic paradigm that are hard-to-grasp. This may be due, on the one hand, to the heterogeneity of the interventions included within this theoretical model, and on the other, to the difficulty in measuring health outcomes.

Understanding the achievements of interventions based on the salutogenic theory could lead to stronger support for this type of actions in the field of public health and the implementation of health and well-being policies.

The main objective of the present review is to explore the type of interventions developed using salutogenic theoretical concept and the effects on physical morbidity, mortality, mental health, health related quality of life (HRQoL) and subjective well-being.

Method

A scoping review¹⁵ was conducted in 2017. Four databases were searched: PubMed, Embase, Web of Science and Scopus, and hand searches through snowballing was performed. We included articles published over the last 10 years (2007/2016), using the following descriptors related to salutogenic models and selected by the authors: (“Evaluation” OR “Program Evaluation” OR “Outcome Assessment”) AND (“Salutogenesis” OR “social capital” OR “health asset” OR “urbanism” OR “sense of coherence” OR “health assets” OR “health assets model” OR “Salutogenic” OR “Salutogenetic” OR “Salutogenic approach” OR “Salutogenic model”) AND (“Mortality” OR “Morbidity” OR “Quality of life” OR “life expectancy” OR “Mental Health”).

This search strategy was adopted to find papers related to salutogenic theory with an asset based approach. The final search strategy needed to be kept broad in order to capture all available information with an asset based approach at individual (SOC) and community level (social capital, urbanism).

We included original articles, reviews, descriptive studies, quasi-experimental and clinical trial articles written in English, French and Spanish. We only took into account interventions that reported evaluations of health outcomes (morbidity, mortality, mental health and/or quality of life). We excluded editorials, letters to the editor, descriptions of experiences without assessments of results, or theoretical dissertations. No other restriction was made.

The database search was carried out in 2017. After removing duplicates, two researchers independently screened the abstracts to identify those which met the eligibility criteria. A random sample of 38 papers was selected to assess the level of agreement between the researchers regarding the classification and extraction of information: a high level of consistency was found (kappa index = 86.4%).

The following information was extracted from each of the original selected articles: authors, year of publication, study design,¹⁶ country, objective, population, duration, variables, results, adverse effects, limitations and biases as presented by the authors themselves.

After an initial reading to become familiar with the contents, common characteristics regarding the focus of interventions and the target population were identified grouping them into four larger groups:

- Individual interventions: encompassing different studies where the intervention was centred on individuals, fundamentally carried out through different psychotherapeutic treatments and with the purpose of modifying cognitive or behavioural aspects.
- Group interventions: interventions in groups of people, sharing a similar health issue tackled through the intervention. The purpose of these interventions is, usually, the modification of cognitive or behavioural aspects.

- Mixed interventions: interventions combining individual and group approaches.
- Intersectoral interventions: interventions aimed at groups or communities to intervene within the environment in which the health problems take place and develop. These interventions include a minimum of two sectoral areas (urbanism, social services, education, health system. . .).

The extracted information was presented by grouping the articles according to the type of intervention and describing the effects relating to health outcomes. To synthesize the information, we performed a descriptive statistical analysis, including frequencies and means calculated. To study mean differences, a student t-test was performed.

Results

Main characteristics of the studies

A total of 669 articles were located in the consulted databases distributed as follows: 115 in PubMed, 175 in Web of Science, 164 in Embase, and 215 in Scopus. Complementary papers potentially eligible were included, a search through snowballing was performed, and seven studies were identified (Fig. 1).

After screening titles and abstracts, 72 papers were retrieved full-text for further reading. A total of 61 articles fulfilled the

inclusion criteria and were finally included in the scoping review. Most part of the studies, 62.29% (38), were published between 2013–2016. The studies included were carried out in 23 countries within the five continents. Were done with women 8.2% (5) of the studies, 3.3% (2) with men, and the remaining 88.5% (54) with men and women. With regard to the average number of months of the monitoring concerning the studies identified, intersectoral interventions were those with the highest monitoring data (86.6 ± 92.9), followed by the individual interventions (20.3 ± 18.1), the group interventions (16.8 ± 22.9), and finally the mixed interventions (13.53 ± 7.1). The biggest category was that of group interventions (39.34% of the studies), followed by intersectoral interventions (34.42%), mixed interventions (14.75%) and individual interventions (11.47%). The main characteristics are described in Table 1.

Quasi-experimental designs were used in 40.98% of the papers, notably including community trials and pre-test/post-test studies. The randomized clinical trials represented 19.67% of the studied sample. Only a total of 4.92% of the interventions were analyzed using qualitative research designs. Important differences existed in the design typology depending on the type of interventions.

The most prevalent methodological designs in each type of interventions were randomized controlled trials for individual interventions (57.14%), quasiexperimental designs for group (41.67%) and mixed interventions (77.68%), and finally descriptive studies for intersectoral interventions (33.33%).




Figure 1. Diagram of the scoping review realized.

Table 1
General characteristic of the studies.

	Type of intervention				
	Individual (n = 7)	Group (n = 24)	Mixed (n = 9)	Intersectoral (n = 21)	Total (n = 61)
<i>Year of publication</i>					
2007–2009	2 (28.57%)	4 (16.67%)	3 (33.33%)	3 (14.29%)	12 (19.67%)
2010–2012	-	3 (12.5%)	1 (11.11%)	7 (33.33%)	11 (18.03%)
2013–2016	5 (71.43%)	17 (70.83%)	5 (55.56%)	11 (52.38%)	38 (62.29%)
<i>Design</i>					
Descriptive (n = 9019 ^a)	-	4 (16.67%)	-	7 (33.33%)	11 (18.03%)
Analytic (n = 2129)	1 (14.29%)	1 (4.17%)	2 (22.22%)	1 (4.76%)	5 (8.20%)
Quasiexperimental (n = 15.142)	2 (28.57%)	10 (41.67%)	7 (77.78%)	6 (28.57%)	25 (40.98%)
RCT (1905)	4 (57.14%)	7 (29.17%)	-	1 (4.76%)	12 (19.67%)
SR/SR (7003 ^b)	-	1 (4.17%)	-	4 (19.05%)	5 (8.20%)
Qualitative (n = 53)	-	1 (4.17%)	-	2 (9.52%)	3 (4.92%)
<i>Monitoring average (months)</i>	20.3 ± 18.1	16.8 ± 22.9	13.53 ± 7.1	86.6 ± 92.9	39.9 ± 64.2
<i>Total size</i>	n = 1055	n = 10.730	n = 11.888	n = 11.578 ^c	n = 35.251
<i>Sample size</i>					
< 101	4 (57.14%)	14 (58.33%)	6 (66.67%)	6 (28.57%)	26 (42.62%)
101–500	3 (42.86%)	6 (25%)	2 (22.22%)	2 (9.52%)	14 (22.95%)
501–1000	-	3 (12.50%)	-	-	6 (9.83%)
> 1000	-	1 (4.17%)	1 (11.11%)	4 (19.04%)	6 (9.83%)
No available	-	-	-	9 (42.85%)	9 (14.75%)

RCT: randomized controlled trial; SR/SR: scoping review/systematic review.

^a 45.4% (5) of the articles did not specify the sample size adopted in the intervention.

^b 80% (4) of the articles did not specify the sample size adopted in the intervention.

^c 42.8% (9) of the articles did not specify the sample size adopted in the intervention.

Type of salutogenic interventions

Table 2 shows the different types of interventions. Most of the individual interventions are focused on health education activities, counselling and/or psychotherapeutic approaches from different approaches (cognitive-behavioural therapy, psychodynamic, occupational therapy).

Group interventions include health education activities, as well as different types of psychotherapy (cognitive-behavioural, psychodynamic, holistic, community). Unlike individual interventions, they are developed in groups of people sharing a similar health problem.

Mixed interventions incorporate combined actions at individual and group level.

Intersectoral interventions incorporate actions aimed at intervening not only on people and communities, but also within the environment in which the health problems take place. These interventions are carried out with the participation of stakeholders from two or more different background areas (urban renewal programs, social participation, governance...).

Relationship between salutogenic interventions and health outcomes

For each intervention type, a summary of the salutogenic interventions is given, describing the main health outcomes improvements and limitations identified (Table 2).

As can be seen in Table 2, studies in the field of individual salutogenic interventions are developed in relation to physical morbidity (human immunodeficiency virus [HIV]), mental health and perceived quality of life and health. These types of interventions led to a variety of health outcomes improvements, notably the following: a decrease in HIV rates, a reduction in anxiety-related symptomatology, an increase in perception of quality of life, higher personal satisfaction, a reduction in psychopathological symptomatology, and a drop in eating disorders.

Group interventions targeted different health topics, ranging from the management of chronic diseases, to mental health and

subjective well-being. Additionally, some papers examined different social dimensions relating to social trust, social capital and social compromise. In these types of interventions, improvements were identified in different fields of study, such as: a reduction in disease prevalence and morbidity burden, improvements in perceived health, greater well-being and perceived quality of life, a decrease in psychopathological symptomatology, as well as other symptoms relating to anxiety and depression, improved functional capacity, lower burnout and a reduction in stress symptomatology.

The selected mixed interventions presented a wide range of topics linked to management of chronic pain, functional capacity improvement, HRQoL, drug consumption, and psychosomatic disorders. These studies identified improvements in different fields of action linked to a reduction in pain-related symptomatology, a drop in insomnia and an increase in dream quality, improvements in HRQoL, an increase of the functional capacity, a decrease of kinesiophobia, higher psychological, physical and social well-being, and greater physical and mental health.

The selected studies in intersectoral interventions identified were focused in urban development, the implementation of community interventions, self-care programmes, social prescription, development of public policies and basic services, governance and multisectorality, health behaviours approach, social capital, and mental health. These interventions led to improvements in HRQoL, an increase in perceived well-being and self-rated health, greater mental health, a reduction in depressive symptomatology, a decrease in preventable mortality for diabetes mellitus, influenza, heart disease and infant mortality.

Positive health outcomes and methodological limitations of the salutogenic interventions

There were important differences between the positive health outcomes depending on the typology of intervention, but most of the studies (85.25%) showed positive findings. None of the studies reported adverse effects.

Two of the identified articles developed by Ciccone et al. in 2014,¹⁷ and Mays et al. in 2016,¹⁸ showed a relation

Table 2
Effects and limitations identified by typology of intervention.

	Individual interventions	Group Interventions	Mixed interventions	Intersectoral interventions
Positive effects	71.43%	87.50%	100.00%	80.95%
Adverse effects	0%	0%	0%	0%
Health outcomes improvements	HIV rate ²²	Prevalence and morbidity burden ^{27,28}	Functional capacity ⁴⁰	Life expectancy at birth, life expectancy in good health, and mortality for a variety of reasons: maternal, infantile and famine mortality ¹⁷
	Anxiety-related symptomatology ²³ Eating disorders ²⁴	Perceived health ^{29,30} Well-being and perceived quality of life ^{31–37}	Pain-related symptomatology ^{40,41} Dream quality ^{40,42,43}	Perceived well-being and self-rated health ^{17,49} Preventable mortality for diabetes mellitus, influenza, heart disease and infant mortality ¹⁸
	Perception of quality of life ^{24,25}	Psychopathological symptomatology, anxiety and depression ^{32,39}	HRQoL ^{40–42,44,45}	HRQoL ^{50–53}
	Personal satisfaction ²⁶ Psychopathological symptomatology ^{24,26}	Stress symptomatology ³⁴ Functional capacity ³⁸ Burnout ³⁹	Kinesophobia ^{40,42} Psychological, physical and social well-being ⁴³ Physical and mental health ^{44,46–48}	Mental health ^{54–56} Depressive symptomatology ⁵⁷
<i>Type of methodological limitations (identified by authors in their own work)</i>				
Sample size	57.14%	37.50%	55.56%	14.29%
Generalization results	28.57%	29.17%	0.00%	28.57%
Selection bias	14.29%	37.50%	33.33%	42.86%
Unvalidated tools	0.00%	0.00%	22.22%	9.52%
Confounding variables control	28.57%	20.83%	11.11%	33.33%
Sample loss	14.29%	16.67%	11.11%	4.76%
Equivalent control group	14.29%	16.67%	0.00%	9.52%

HIV: human immunodeficiency virus; HRQoL: health related quality of life.

between interventions and mortality reduction. Both studies are intersectoral interventions and they were based on the influence of health policies and governance mechanisms and their impact on health.

The study of Mays et al.¹⁸ aimed to measure the extent and nature of multisector contributions to population health activities and how these contributions affect community mortality rates. The study of Ciccone et al.¹⁷ about governance mechanisms and health outcomes in low and middle income countries suggest that governance mechanism may influence health outcomes in different settings (life expectancy at birth, life expectancy in good health, as well as reduced mortality).

The mixed interventions are those that present the highest degree of positive effects (100%). The positive effects identified in this review did not take into account the methodological design of each study, which would have had an impact on the findings discussed. Significant methodological limitations were found by the authors of the papers (75.41%). The most common biases identified were associated to participant selection bias in 36% of the cases, followed by the sample size (34.43%). To a lesser extent, the absence of a control group or a lack of equivalence in the control group were reported as potential limitations (11.48%), important sample losses occurred during the implementation of the studies (11.48%) and in 6.56% of the studies the use of non-validated tools was discussed as a potential limitation.

Conclusions

This research has offered an overview of interventions adopting a salutogenic framework, from the individual to the intersectoral level, synthesizing the variety of actions which can be included under this paradigm, and describing their impacts on health and their study limitations.

Recently, the number of studies carried out in this field of study has significantly increased. Most of the studies described positive effects on health, especially in the field of mental health. Grouped,

mixed or intersectoral interventions showed better health outcomes than de individual ones. This may be due to the incorporation of a more collective approach to the problem. Such an approach may enhance social support and social networks, thus addressing a broader range of health determinants and potentially increasing better health outcomes.¹⁴ The majority of positive effects are related to perceived quality of life and health, followed by mental health and physical morbidity reduction.

Most of the interventions reflect the salutogenic theories only partially, through incorporating some of their elements, such as SOC or GRRs which were the focus of this review. A systemic salutogenic approach, that seeks to act not only on the individuals but also on the communities and contexts in which they live, could have an expanded effect in terms of improving the health outcomes of the population, as seen in the positive impact associated with some of the intersectoral interventions.^{17,18} However, these positive results could depend on the fact that longer periods of monitoring are present in intersectoral actions

Despite the fact that different authors^{1–3,19,20} have tried to define the salutogenic theory it is necessary to reach a consensus that defines the minimal characteristics that a salutogenic intervention should follow and validate a tool which evaluates the salutogenic contents of the interventions. Reviews of salutogenic-based interventions and their effects on population health are limited.

Our research has some methodological limitations which can affect the interpretation as well as the generalization of the results. Some of the terms used such as health assets, do not have normalized descriptors in the databases, thus affecting the sensitivity of the search process.²¹ The search strategy was limited at identifying articles related to the application of the salutogenic paradigm in health interventions, thus eventually excluding other studies adopting the paradigm in other sectors and studies using this approach without qualifying them as salutogenic. On the other hand, the use of standardized descriptors in the search strategy of our study has facilitated the inclusion of papers that, in other

contexts, would not meet the minimum criteria to consider them as salutogenic.

In some cases, data extraction was challenging, since not all studies provided detailed information on their methodological design. In some studies, the dependent and independent variables were not clearly identified and the authors of the present work had to classify them according to the available information.

As for the limitation identified in the included papers these studies need to be conducted with improved monitoring and evaluation designs. Future research should centre on developing a typology of salutogenic interventions, potential beneficiaries and effects on health. Tools as well as indicators are also necessary to better understand the impact of those interventions on the population health.

Methodological rigour must be established; studies should be carried out under stricter conditions which allow confirming the efficacy of interventions. Furthermore, additional tracking should be encouraged to account for the medium and long-term effects of salutogenic programmes.

Qualitative research in this area can support in the understanding of the mechanisms that explain the health improvement, including the range of outcomes identified by the population itself.

Further research may also need to account for a potential publication bias in the sense of a tendency to publish only articles with positive results which could explain the high rate of positive effects identified in the studies included in this scoping review.

The lack of clinical trials, the wide variety of methodological designs and the scientific validity of some of the interventions could explain the difficulties for the practical implementation of this theoretical model. Nonetheless, most of the findings of the studies included in this review seem to suggest that salutogenic interventions can achieve positive effects and that could have numerous empirical applications.

We hope the results obtained in this review will be useful for other researchers, especially in terms of understanding the salutogenic theoretical concept and the range of interventions types and health outcomes that it can achieved as described by the authors in the scientific literature.

Editor in charge

Clara Bermúdez Tamayo.

Authorship contributions

O. Suárez Álvarez, C. Álvarez-Dardet and M.T. Ruiz Cantero made the design of the scoping review. O. Suárez Álvarez made the data collection and classification of the information. R. Cofiño made the concordance analyses with O. Suárez Álvarez. R. Cofiño, C. Álvarez-Dardet, V. Cassetti, O. Suárez Álvarez and M.T. Ruiz Cantero made the analysis and interpretation of the results write the drafting article and the final approval of the version to be published.

Acknowledgements

The authors wish to thank to Floriane Nguyen, Gwendolijn Boonekamp, Patricia Perez Wilson and Jorge Marcos for their comments on an earlier draft and for her help preparing this manuscript. This paper is part of the PhD training of Oscar Suárez in the University of Alicante Health Sciences Programme and will be used as part of his PhD dissertation as compendium of articles.

Funding

None.

Conflicts of interest

Two of the authors (C.A.D. and M.T.R.C.) belongs to the GAC-ETA SANITARIA editorial committee, but were not involved in the editorial process of the manuscript.

Bibliografía

- Antonovsky A, Maoy B, Dowty N, et al. Twenty-five years later: a limited study of the sequelae of the concentration camp experience. *Soc Psychiatry Psychiatr Epidemiol.* 1971;6:186–93.
- Antonovsky A. *Unraveling the mystery of health. How people manage stress and stay well.* San Francisco: Jossey-Bass; 1987. p. 238.
- Antonovsky A. The salutogenic model as a theory to guide health promotion. *Health Promot Int.* 1996;11:11–8.
- Becker CM, Glascoff MA, Felts M. Salutogenesis 30 years later: where do we go from here? *International Electronic Journal of Health Education.* 2010;13:25–32.
- Vaandrager L, Kennedy L. The application of salutogenesis in communities and neighborhoods. In: Mittelmark MB, Sagy S, Eriksson M, et al, editors. *The handbook of salutogenesis.* Cham (CH): Springer; 2017. p. 159–70.
- Eriksson M, Lindström B, Lilja J. A sense of coherence and health Salutogenesis in a societal context: Aland, a special case? *J Epidemiol Community Health.* 2007;61:684–8.
- Bauer GF, Roy M, Bakibinga P, et al. Future directions for the concept of salutogenesis: a position article. *Health Promot Int.* 2019, pii: daz057.
- Jensen BB, Dür W, Buijs G. The application of salutogenesis in schools. In: Mittelmark MB, Sagy S, Eriksson M, et al, editors. *The handbook of salutogenesis.* Cham (CH): Springer; 2017. p. 225–35.
- Pelikan JM. The application of salutogenesis in healthcare settings. In: Mittelmark MB, Sagy S, Eriksson M, et al, editors. *The handbook of salutogenesis.* Cham (CH): Springer; 2017. p. 261–6.
- Taylor JS. Salutogenesis as a framework for child protection: literature review. *J Adv Nurs.* 2004;45:633–43.
- Aujoulat I, Mustin L, Martin F, et al. The application of salutogenesis to health development in youth with chronic conditions. In: Mittelmark MB, Sagy S, Eriksson M, et al, editors. *The handbook of salutogenesis.* Cham (CH): Springer; 2017. p. 337–44.
- Eriksson M, Lindström B. Antonovsky's sense of coherence scale and the relation with health: a systematic review. *J Epidemiol Community Health.* 2006;60:376–81.
- González Hernández MD, Santos Sancho JM. directors. *La influencia de la salud percibida en la mortalidad de las personas mayores.* [Tesis doctoral]. Madrid: Universidad Complutense de Madrid; 2017.
- Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. *PLoS Med.* 2010;7:e1000316.
- Peters M, Godfrey C, McInerney P, et al. *The Joanna Briggs Institute Reviewers' Manual 2015: methodology for JBI scoping reviews.* Adelaide, SA, Australia: The Joanna Briggs Institute; 2015. (Accessed 07/06/2016.) Available at: <http://joannabriggs.org/assets/docs/sumari/Reviewers-Manual.Methodology-for-JBI-ScopingReviews.2015.v2.pdf>.
- Burgos R, Chicharro JA, Bobenrieth M. *Metodología de investigación y escritura científica en clínica.* Granada: Escuela Andaluza de Salud Pública; 1994. p. 457.
- Ciccione DK, Vian T, Maurer L, et al. Linking governance mechanisms to health outcomes: a review of the literature in low- and middle-income countries. *Soc Sci Med.* 2014;117:86–95.
- Mays GP, Mamaril CB, Timsina LR. Preventable death rates fell where communities expanded population health activities through multisector networks. *Health Affairs.* 2016;35:2005–13.
- Lindstrom B, Eriksson M. The salutogenic approach to the making of HiAP/healthy public policy: illustrated by a case study. *Glob Health Promot.* 2009;16:17–28.
- Rivera de los Santos F, Ramos Valverde P, Moreno Rodríguez C, et al. Salutogenic model analysis in Spain: application in public health and implications for asset health model. *Rev Esp Salud Publica.* 2011;85:129–39.
- Van Bortel T, Wickramasinghe ND, Morgan A, et al. Health assets in a global context: a systematic review of the literature. *BMJ Open.* 2019;9:e023810.
- Drame FM, Crawford EE, Diouf D, et al. A pilot cohort study to assess the feasibility of HIV prevention science research among men who have sex with men in Dakar Senegal. *J Int AIDS Soc.* 2013;16 Suppl 3:18753.
- Jensen JF, Egerod I, Bestle MH, et al. A recovery program to improve quality of life, sense of coherence and psychological health in ICU survivors: a multicenter randomized controlled trial, the RAPIT study. *Intensive Care Med.* 2016;42:1733–43.
- Dingemans AE, Danner UN, Donker JM, et al. The effectiveness of cognitive remediation therapy in patients with a severe or enduring eating disorder: a randomized controlled trial. *Psychother Psychosom.* 2013;83:29–36.
- Knekt P, Heinonen E, Härkäpää K, et al. The Helsinki Psychotherapy Study Group Randomized trial on the effectiveness of long- and short-term psychotherapy on psychosocial functioning and quality of life during a 5-year follow-up. *Psychiatry Res.* 2015;229:381–8.
- Gunnarsson AB, Eklund M. The tree theme method as an intervention in psychosocial occupational therapy: client acceptability and outcomes. *Aust Occup Ther J.* 2009;56:167–76.

27. Stenberg U, Haaland-Overby M, Fredriksen K, et al. A scoping review of the literature on benefits and challenges of participating in patient education programs aimed at promoting self-management for people living with chronic illness. *Patient Educ Couns.* 2016;99:1759–71.
28. Struck P, Ventegodt S. Clinical holistic medicine: teaching orgasm for females with chronic anorgasmia using the Betty Dodson method. *ScientificWorldJournal.* 2008;8:883–95.
29. de Souza EM, Grundy E. Intergenerational interaction, social capital and health: results from a randomised controlled trial in Brazil. *Soc Sci Med.* 2007;65:1397–409.
30. Im H, Rosenberg R. Building social capital through a Peer-Led Community Health Workshop: a pilot with the Bhutanese refugee community. *J Community Health.* 2016;41:509–17.
31. Chlond M, Bergmann F, Gütthlin C, et al. Patient education for patients with Parkinson's disease: a randomised controlled trial. *Basal Ganglia.* 2016;6:25–30.
32. Schrank B, Brownell T, Jakaite Z, et al. Evaluation of a positive psychotherapy group intervention for people with psychosis: pilot randomised controlled trial. *Epidemiol Psychiatr Sci.* 2015;25:1–12.
33. Calandri E, Graziano F, Borghi M, et al. Improving the quality of life and psychological well-being of recently diagnosed multiple sclerosis patients: preliminary evaluation of a group-based cognitive behavioral intervention. *Disabil Rehabil.* 2017;39:1474–81.
34. Esch T, Sonntag U, Esch SM, et al. Stress management and mind-body medicine: a randomized controlled longitudinal evaluation of students' health and effects of a behavioural group intervention at a middle-size German University (SM-MESH). *Forschende Komplementarmedizin.* 2013;20:129–37.
35. Graziano F, Calandri E, Borghi M, et al. The effects of a group-based cognitive behavioral therapy on people with multiple sclerosis: a randomized controlled trial. *Clin Rehabil.* 2014;28:264–74.
36. Ko HC, Kuo FY. Can blogging enhance subjective well-being through self-disclosure? *Cyberpsychol Behav.* 2009;12:75–9.
37. Strange C, Bremner A, Fisher C, et al. Mothers' group participation: associations with social capital, social support and mental well-being. *J Adv Nurs.* 2016;72:85–98.
38. Sakurai R, Yasunaga M, Murayama Y, et al. Long-term effects of an intergenerational program on functional capacity in older adults: results from a seven-year follow-up of the REPRINTS study. *Arch Gerontol Geriatr.* 2016;64:13–20.
39. Wild K, Scholz M, Ropohl A, et al. Strategies against burnout and anxiety in medical education – implementation and evaluation of a new course on relaxation techniques (Relacs) for medical students. *PLoS One.* 2014;9:e114967.
40. Hållstam A, Löfgren M, Benson L, et al. Assessment and treatment at a pain clinic: a one-year follow-up of patients with chronic pain. *Scand J Pain.* 2016;10:36–42.
41. Bergström M, Ejelöv M, Mattsson M, et al. One-year follow-up of body awareness and perceived health after participating in a multimodal pain rehabilitation programme – a pilot study. *Eur J Physiother.* 2014;16:246–54.
42. Hållstam A, Löfgren M, Svensén C, et al. Patients with chronic pain: one-year follow-up of a multimodal rehabilitation programme at a pain clinic. *Scand J Pain.* 2016;10:36–42.
43. Lev-ari S, Zilcha-Mano S, Rivo L, et al. A prospective pilot clinical trial of “The work” meditation technique for survivors of breast cancer. *European Journal of Integrative Medicine.* 2013;5:487–94.
44. Ventegodt S, Kandel I, Merrick J. Clinical holistic medicine in the recovery of working ability. A study using Antonovsky salutogenesis. *International Journal on Disability and Human Development.* 2008;7:219–22.
45. Fernros L, Furhoff AK, Wändell PE. Improving quality of life using compound mind-body therapies: evaluation of a course intervention with body movement and breath therapy, guided imagery, chakra experiencing and mindfulness meditation. *Qual Life Res.* 2008;17:367–76.
46. Ventegodt S, Thegler S, Andreassen T, et al. Clinical holistic medicine (mindful, short-term psychodynamic psychotherapy complemented with bodywork) in the treatment of experienced physical illness and chronic pain. *ScientificWorldJournal.* 7, 310–6.
47. Ventegodt S, Thegler S, Andreassen T, et al. Self-reported low self-esteem. Intervention and follow-up in a clinical setting. *ScientificWorldJournal.* 2007;7:299–305.
48. Schori M, Sapir Y, Lawental E. Long-term residential substance abuse treatment for women: lessons learned from Israel. *Subst Abuse Rehabil.* 2012;3:71–81.
49. Ichida Y, Hirai H, Kondo K, et al. Does social participation improve self-rated health in the older population? A quasi-experimental intervention study. *Soc Sci Med.* 2013;94:83–90.
50. Tan KK, Chan SWC, Wang W, et al. A salutogenic program to enhance sense of coherence and quality of life for older people in the community: a feasibility randomized controlled trial and process evaluation. *Patient Educ Couns.* 2016;99:108–16.
51. Bungay H, Clift S. Arts on prescription: a review of practice in the U.K. *Perspect Public Health.* 2010;130:277–81.
52. Møller V. Quality of life in South Africa – the first ten years of democracy. *Social Indicators Research.* 2007;81:181–201.
53. Parsfield M, Morris DM, Bola DM, et al. Report: Community Capital – The value of connected communities. London: Royal Society for the encouragement of Arts, Manufactures and Commerce; 2015. Available at: www.thersa.org/discover/publications-and-articles/reports/community-capital-the-value-of-connected-communities/Download.
54. Hadley C, Stevenson EGJ, Tadesse Y, et al. Rapidly rising food prices and the experience of food insecurity in urban Ethiopia: impacts on health and well-being. *Soc Sci Med.* 2012;75:2412–9.
55. Houle JN. Mental health in the foreclosure crisis. *Soc Sci Med.* 2014;118:1–8.
56. Wilson RL. Mental health recovery and quilting: evaluation of a grass-roots project in a small, rural, Australian Christian church. *Issues Ment Health Nurs.* 2014;35:292–8.
57. Coll-Planas L, Del Valle Gómez G, Bonilla P, et al. Promoting social capital to alleviate loneliness and improve health among older people in Spain. *Health Soc Care Community.* 2015;25:145–7.