

Original

Immigration status and depression in Spain: analysis of the European Health Interview Surveys of 2014 and 2020

Luisa N. Borrell^{a,b,*}, Julia Díez^b, Sara Yago-González^b, Nerea Lanborena^{c,d}, Elena Rodríguez-Álvarez^{c,d}^a Department of Epidemiology and Biostatistics, Graduate School of Public Health and Health Policy, The City University of New York, New York, NY, United States^b Public Health and Epidemiology Research Group, Department of Surgery, Medical and Social Sciences, Faculty of Medicine and Health Sciences, Universidad de Alcalá, Alcalá de Henares, Madrid, Spain^c Department of Nursing I, University of the Basque Country, Leioa, Bizkaia, Spain^d OPIK-Research Group for Social Determinants of Health and Demographic Change, University of the Basque Country, Leioa, Bizkaia, Spain

ARTICLE INFO

Article history:

Received 21 October 2024

Accepted 21 December 2024

Keywords:

Immigrants

Depression

Social inequalities

Spain

ABSTRACT

Objective: We examined the association between immigration status and depression in a nationally representative sample of adults in Spain. In addition, we assessed whether this association differed by sex/gender and social support.

Method: We used de-identified data from the European Health Interview Survey conducted in Spain in 2014 (n=21,226) and 2020 (n=20,136). Our study outcomes were self-reported diagnosis of depression and antidepressant use. We fitted Poisson regression models to quantify the association between immigration status and each outcome, before and after adjusting for age, sex/gender, marital status, educational level, employment status, smoking status, healthcare use, social support, and self-rated health. We obtained prevalence ratios (PR) and 95% confidence intervals (95%CI). We also tested interaction terms to evaluate whether the associations of interest differed by survey year and further by sex/gender and with social support.

Results: In 2014 and 2020, we observed a lower prevalence of depression among immigrant adults than among native Spanish adults (adjusted PR₂₀₁₄ = 0.53, 95%CI: 0.40- 0.70; PR₂₀₂₀ = 0.71, 95%CI: 0.55-0.91). However, for antidepressant use, this association was significant only in 2014 (adjusted PR₂₀₁₄ = 0.36, 95%CI: 0.24-0.55). Although the association of immigration status with antidepressant use differed with survey year, these associations did not change with sex/gender or social support.

Conclusions: Our findings call attention to depression and mental health-related outcomes in Spain, regardless of immigration status, sex/gender or social support.

© 2024 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Estado migratorio y depresión en España: análisis de las Encuestas Europeas de Salud de 2014 y 2020

RESUMEN

Palabras clave:

Inmigrantes

Depresión

Desigualdades sociales

España

Objetivo: Examinamos la asociación entre el estado migratorio y la depresión en una muestra representativa de ámbito nacional de adultos en España. Además, evaluamos si esta asociación difiere según el sexo/género y el apoyo social.

Método: Utilizamos datos anonimizados de la Encuesta Europea de Salud realizada en España en 2014 (n = 21.226) y en 2020 (n = 20.136). Los resultados de nuestro estudio fueron el diagnóstico autoinformado de depresión y el uso de antidepresivos. Usamos modelos de regresión de Poisson para cuantificar la asociación entre el estado migratorio y cada resultado, antes y después de ajustar por edad, sexo/género, estado civil, nivel educativo, situación laboral, tabaquismo, uso de atención médica, apoyo social y salud autopercebida. Obtuvimos razones de prevalencia (RP) e intervalos de confianza del 95% (IC95%). También examinamos términos de interacción para evaluar si las asociaciones de interés diferían según el año de la encuesta, y además según el sexo/género y el apoyo social.

Resultados: En 2014 y 2020 observamos una menor prevalencia de depresión en los inmigrantes que en los adultos nacidos en España (RP₂₀₁₄ ajustada = 0,53, IC95%: 0,40-0,70; RP₂₀₂₀ = 0,71, IC95%: 0,55-0,91). Sin embargo, para el uso de antidepresivos esta asociación solo fue significativa en 2014 (RP₂₀₁₄ ajustada = 0,36, IC95%: 0,24-0,55). Aunque la asociación del estado migratorio con el uso de antidepresivos difiere según el año de la encuesta, estas asociaciones no cambiaron con el sexo/género o el apoyo social.

* Corresponding author.

E-mail address: Luisa.Borrell@sph.cuny.edu (L.N. Borrell).

Conclusiones: Nuestros hallazgos llaman la atención sobre la depresión y las consecuencias relacionadas con la salud mental en España, independientemente del estado migratorio, el sexo/género y el apoyo social.

© 2024 SESPAS. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

In Europe, previous studies have reported that immigrants generally exhibit a higher prevalence of depressive symptoms than native populations.^{1–4} In Spain, a country with the third highest proportion of immigrant population in the European Union,⁵ evidence for this association is mixed,^{6–9} with studies reporting a higher prevalence,⁹ lower prevalence,^{6,7} or no difference in the prevalence of depression between immigrants and native Spanish adults.⁸ In addition, immigrants are less likely to consume antidepressants.^{6,8} Thus, it is important to examine the associations of immigration status with depression and related outcomes.

Social, economic, and psychosocial factors influenced the association between immigration status and depression.^{1,10,11} Factors contributing to these social inequalities include unemployment, lower social support, and previous mental health conditions. Precarious working conditions, for example, have been shown to significantly increase the risk of depression among immigrant workers.^{12,13} Additionally, immigrants often face unique challenges such as discrimination, which is linked to depressive symptoms.¹⁴ Gender inequalities have also been documented among immigrants, with depression being more common among women than men.⁸ Similarly, social support tends to buffer the stress related to immigration status,¹⁵ and thus, may change the relationships of immigration status with depression and antidepressant use.

Spain presents unique challenges regarding the migration processes, as noted in previous studies.^{16,17} Currently, 17.1% of the population has an immigrant background, mostly from South America or Africa.⁵ However, our understanding of social inequalities in mental health, based on immigration status, remains limited. Therefore, we aimed to examine whether there is an association of immigration status with depression and antidepressant use in adults who participated in the European Health Interview Survey (EHIS) of Spain in 2014 and 2020, and whether this association differs by sex/gender and social support.

Method

Study design and data collection

We conducted a cross-sectional analysis of immigrant and native Spanish adults aged 18 or older who participated in the EHIS conducted in Spain in 2014 and 2020. These surveys used a stratified multistage sampling approach to ensure the representation of Spain's noninstitutionalized population. Data collection for the 2014 survey took place between February 2014 and January 2015, while the 2020 survey was conducted between July 2019 and July 2020. The response rates for the two surveys were 71% and 59%, respectively.

Study variables

Consistent with a previous study in Spain,¹⁸ we assessed depression using self-reported information about two survey questions: "Have you ever been depressed?" and "Has a medical doctor told you that you had depression". We deemed a positive response to both questions as indicative of depression. Additionally, we consid-

ered the use of antidepressants prescribed in the last two weeks if individuals reported depression. We measured immigration status by country of birth: born in Spain or outside Spain, including adults born in countries with a human development index (HDI) <0.80.¹⁹ The cutoff point is recommended for comparison purposes, as countries with HDI ≥0.80 have social conditions similar to Spain, and thus, immigrants from these countries may have similar outcomes to the Spanish population. In addition, we used country of birth to determine the continent of origin of the immigrant population as follows: Asia, Latin America, Europe, and Africa.

Consistent with previous studies,^{6,12,20–23} we considered age (continuous and categorical), sex/gender (men or women), marital status (married or not), educational level (no formal education or primary, secondary, or tertiary), employment status, occupation or social class, use of primary healthcare (last visit to a general practitioner in the past four weeks, between four weeks and 12 months, and 12 months or more/never), social support, smoking status, and self-rated health.

Employment status was specified using information on contractual conditions and the actual employment situation as follows: permanent as someone who had a permanent contract and was currently working; precarious as someone who was currently working part-time fixed-term contracts, or with informal employment arrangements, including unspecified situations and unpaid family workers; business as someone who was working as a business or professional employee or worker independent; others included individuals who were working because they retired or pre-retired, students, or a person with disabilities; and unemployed included individuals who were not working but had a previous occupation. Occupation or social class (hereafter, social class) was collected using the following categories: directors of establishments with ten or more employees, self-employed or intermediate occupations, supervisors/technical occupations, primary sector/semi-qualified occupations, and non-qualified occupations. For analytical purposes, we specified social class as non-manual (including the first three categories) or manual (including the last two categories).

We used questions from the social support scale Oslo³²² regarding the number of people one can count on, the extent to which people care about what happened to people, or people can get help from neighbors. The sum of the responses to these questions ranged from 3 to 14 points. We categorized social support as low (3–8), moderate (9–11), or strong (12–14). Responses to the question "Do you currently smoke?" (yes, every day; yes, but not daily; no, currently but I have smoked; and I have never smoked) to specify smoking status as current (participants responded that they currently smoke daily or not), former (those who have smoked but not currently), or never (those who have never smoked). Self-rated health was collected via the question "In the last 12 months would you say that your health has been very good, good, regular, bad, or very bad?". We used categorized self-rated health as very good/good (very good/good) and fair/poor (regular, bad, or very bad).^{24–26} All variables were self-reported.

Study population

Of the total sample (n = 22,842 for 2014 and n = 22,072 for 2020), we excluded records of individuals who were < 18 years of age at

Table 1
Descriptive statistics of the study population by immigration status. European Health Interview Surveys in Spain, 2014 and 2020.

Variable, % (SE)	2014			2020		
	Immigrant, n = 1,378 10.2 (0.32)	Native, n = 19,848 89.8 (0.32)	Total, n = 21,226	Immigrant, n = 1,506 12.3 (0.34)	Native, n = 18,630 87.7 (0.34)	Total, n = 20,136
<i>Continent of origin</i>						
Asia	4.6 (0.68)			6.4 (0.89)		
Latin America	48.8 (1.70)			48.6 (1.52)		
Europe	21.3 (1.38)			21.4 (1.25)		
Africa	25.2 (1.57)			23.7 (1.29)		
<i>Age (years), mean (SE)</i>						
18-24	38.4 (0.44)	50.1 (0.16)	48.9 (0.15)	41.0 (0.39)	51.4 (0.17)	50.0 (0.16)
25-34	41.6 (1.72)	22.2 (0.41)	24.2 (0.41)	32.6 (1.48)	20.4 (0.42)	22.0 (0.42)
35-44	30.2 (1.49)	19.8 (0.33)	20.9 (0.33)	30.7 (1.37)	17.5 (0.34)	19.1 (0.34)
45-64	18.7 (1.33)	18.9 (0.34)	18.8 (0.33)	21.7 (1.23)	19.4 (0.36)	19.6 (0.35)
>65	6.5 (0.79)	15.2 (0.29)	14.3 (0.28)	10.2 (0.88)	17.3 (0.33)	16.4 (0.31)
	2.9 (0.59)	23.9 (0.33)	21.8 (0.31)	4.7 (0.60)	25.4 (0.36)	22.8 (0.33)
<i>Sex/gender</i>						
Men	44.6 (1.70)	49.6 (0.43)	49.1 (0.42)	44.5 (1.52)	49.9 (0.46)	49.2 (0.44)
<i>Marital status^a</i>						
Married	61.3 (1.68)	60.4 (0.42)	60.5 (0.42)	59.2 (1.50)	60.1 (0.45)	60.0 (0.44)
<i>Educational level</i>						
No formal/primary	48.5 (1.71)	52.6 (0.43)	52.2 (0.42)	43.1 (1.51)	47.5 (0.46)	46.9 (0.44)
Secondary	37.5 (1.64)	27.6 (0.39)	28.6 (0.39)	42.6 (1.50)	30.9 (0.43)	32.3 (0.42)
Tertiary	14.0 (1.04)	19.8 (0.35)	19.2 (0.33)	14.3 (1.03)	21.6 (0.38)	20.8 (0.36)
<i>Social class</i>						
No manual	18.6 (1.23)	40.2 (0.42)	38.0 (0.41)	18.6 (1.21)	41.3 (0.45)	38.5 (0.43)
<i>Employment status</i>						
Permanent	23.5 (1.36)	27.8 (0.38)	27.4 (0.37)	31.2 (1.41)	33.1 (0.43)	32.9 (0.42)
Precarious	31.6 (1.61)	16.5 (0.35)	18.0 (0.36)	35.6 (1.47)	16.4 (0.37)	18.7 (0.38)
Business	5.4 (0.70)	8.5 (0.70)	8.1 (0.23)	5.2 (0.64)	8.1 (0.25)	7.8 (0.24)
Other	11.6 (1.18)	35.8 (0.40)	33.4 (0.38)	10.5 (0.92)	35.6 (0.42)	32.5 (0.39)
Unemployed	27.9 (1.54)	11.4 (0.28)	13.1 (0.30)	17.6 (1.14)	6.8 (0.24)	8.1 (0.25)
<i>Healthcare use^a</i>						
Within 4 weeks	26.9 (1.57)	29.9 (0.38)	29.6 (0.38)	20.2 (1.18)	21.6 (0.37)	21.4 (0.35)
4 weeks - 1 year	48.2 (1.70)	48.2 (0.43)	48.2 (0.42)	59.8 (1.50)	61.4 (0.45)	61.2 (0.43)
> 1 year/never	24.9 (1.46)	21.9 (0.37)	22.2 (0.36)	20.0 (1.27)	17.0 (0.36)	17.4 (0.35)
<i>Social support</i>						
Low	13.2 (1.07)	4.9 (0.18)	5.7 (0.19)	11.4 (0.92)	3.4 (0.15)	4.4 (0.17)
Moderate	53.6 (1.70)	33.3 (0.40)	35.4 (0.41)	45.9 (1.51)	29.4 (0.42)	31.4 (0.41)
Strong	39.2 (1.62)	61.8 (0.42)	58.9 (0.42)	42.7 (1.51)	67.2 (0.43)	64.2 (0.42)
<i>Smoking status</i>						
Current	23.1 (1.42)	26.2 (0.38)	26.0 (0.37)	18.9 (1.22)	23.4 (0.40)	22.8 (0.38)
Former	17.7 (1.26)	27.7 (0.38)	26.7 (0.36)	13.3 (1.00)	24.6 (0.38)	23.2 (0.36)
Never	59.2 (1.66)	46.0 (0.43)	47.3 (0.42)	67.8 (1.42)	52.0 (0.46)	53.9 (0.44)
<i>Self-rated health</i>						
Fair/poor	26.7 (1.53)	30.3 (0.38)	29.9 (0.38)	22.5 (1.23)	25.4 (0.38)	25.0 (0.36)
<i>Depression, yes</i>						
Depression medications, yes ^a	4.6 (0.68)	9.6 (0.24)	9.1 (0.22)	5.2 (0.63)	7.6 (0.23)	7.3 (0.21)
	2.0 (0.39)	6.2 (0.19)	5.8 (0.18)	3.2 (0.49)	4.9 (0.18)	4.7 (0.17)

SE: standard error.

^a All p-values for chi-squared statistics comparing distribution by immigration status were significant except for marital status and healthcare use in both 2014 and 2020 where p-values were > 0.05.

the time of the interview (n = 521 and 503), records without information on marital status (n = 22 and 64), social support (n = 39 and 26), employment status (n = 61 and 29), smoking (n = 13 and 19), occupation or social class (n = 523 and 934), and records of individuals from countries with an HDI \geq 0.80 (n = 437 and 361). These exclusions yielded final analytical samples of 21,226 and 20,136 in 2014 and 2020, respectively.

Statistical analysis

We calculated descriptive statistics for selected characteristics according to immigration status using mean (\pm standard deviation), frequencies, and percentages, depending on whether the variable was continuous or categorical. We also calculated the prevalence of depression for each characteristic in the total population and by immigration status. We used t-test for continuous variables and chi-square test for categorical variables to determine significant differences or associations, respectively. We fitted a Poisson regression to obtain prevalence ratios (PR) and their 95% confidence intervals (95%CI) and quantify the association of immigration status

with depression and antidepressant use before and after adjusting for age, sex/gender, marital status, educational level, employment status, smoking status, healthcare use, social support, and self-rated health. To determine whether the association varied with the survey year, an interaction term between immigration status and survey year was tested in the adjusted model. Additionally, we examined a three-way interaction to determine whether the association between immigration status and depression varied with sex/gender and with social support in each survey.

All data management and analyses were performed using SAS version 9.4 accounting for the sampling design used in the surveys. Table 1 presents the unweighted sample sizes. However, all estimates presented in the tables were weighted, that is, proportions, standard errors (SE), PR, and 95%CI.

Results

Table 1 displays key characteristics of the study population stratified by immigration status in EHIS 2014 and 2020. Most of the immigrant population originated from Latin American countries.

Table 2

Prevalence estimates of depression for selected characteristics of adults aged 18 years or older by immigration status. European Health Interview Surveys in Spain, 2014 and 2020.

Characteristics	2014		2020	
	Immigrant Yes, % (SE)	Native Yes, % (SE)	Immigrant Yes, % (SE)	Native Yes, % (SE)
<i>Overall</i>	4.6 (0.68)	9.6 (0.24)	5.2 (0.63)	7.6 (0.23)
<i>Continent of origin</i>				
Asia	-		0.8 (0.85)	
Latin America	5.9 (1.12)		6.2 (0.92)	
Europe	4.7 (1.38)		8.1 (1.86)	
Africa	2.7 (1.08)		1.7 (0.78)	
<i>Age, years</i>				
18-34	4.2 (1.21) ^a	2.9 (0.35)	2.7 (0.89)	3.0 (0.42)
25-34	4.1 (1.01)	6.0 (0.46)	4.4 (1.02)	4.5 (0.45)
35-44	6.5 (1.64)	10.0 (0.58)	5.3 (1.18)	7.1 (0.51)
45-64	4.5 (1.95)	13.8 (0.71)	10.4 (2.71)	9.4 (0.57)
65+	3.3 (2.11)	15.8 (0.54)	15.9 (5.45)	12.7 (0.51)
<i>Sex/gender</i>				
Men	1.4 (0.53)	5.7 (0.28)	2.6 (0.69)	5.1 (0.26)
Women	7.1 (1.14)	13.2 (0.38)	7.3 (0.99)	10.1 (0.36)
<i>Marital status</i>				
Married	2.8 (0.64)	9.2 (0.31) ^a	4.4 (0.76) ^a	7.0 (0.28)
Non-married	7.5 (1.42)	10.1 (0.36)	6.3 (1.10)	8.6 (0.37)
<i>Education</i>				
No formal/primary	4.3 (0.88) ^a	13.5 (0.38)	3.7 (0.84) ^a	10.8 (0.38)
Secondary	5.8 (1.36)	6.1 (0.37)	6.0 (1.03)	5.9 (0.39)
Tertiary	2.3 (0.88)	4.1 (0.36)	7.2 (1.93)	3.0 (0.29)
<i>Social class</i>				
No manual	3.1 (1.01) ^a	6.4 (0.30)	5.0 (1.31) ^a	5.7 (0.31)
Manual	4.9 (0.80)	11.8 (0.34)	5.2 (0.72)	8.9 (0.32)
<i>Employment status</i>				
Permanent	3.2 (0.97)	4.4 (0.32)	2.7 (0.77)	4.0 (0.30)
Precarious	4.1 (1.16)	6.3 (0.53)	4.9 (1.06)	4.8 (0.50)
Business	-	4.7 (0.61)	3.5 (1.94)	2.3 (0.43)
Other	6.9 (2.13)	16.2 (0.48)	12.9 (3.19)	13.1 (0.46)
Unemployed	6.2 (1.65)	9.9 (0.76)	6.1 (1.52)	9.8 (0.99)
<i>Healthcare use</i>				
Within 4 weeks	8.5 (1.91)	16.7 (0.55)	13.7 (2.07)	15.0 (0.65)
4 weeks – 1 year	3.9 (0.81)	8.3 (0.32)	3.8 (0.75)	6.4 (0.26)
>1 year/never	1.6 (0.78)	2.8 (0.30)	0.8 (0.59)	2.6 (0.34)
<i>Social support</i>				
Low	6.7 (1.77) ^a	19.3 (1.50)	9.6 (2.32)	15.5 (1.53)
Moderate	4.7 (1.05)	10.5 (0.43)	6.2 (1.03)	9.2 (0.46)
Strong	3.6 (0.91)	8.3 (0.28)	2.9 (0.76)	6.5 (0.26)
<i>Smoking status</i>				
Current	4.9 (1.27) ^a	8.9 (0.48)	7.2 (1.76) ^a	8.1 (0.50) ^a
Former	4.2 (1.44)	8.8 (0.44)	5.7 (1.77)	6.9 (0.43)
Never	4.5 (0.94)	10.5 (0.35)	4.5 (0.72)	7.7 (0.31)
<i>Self-rated health</i>				
Very good/good	1.8 (0.46)	3.4 (0.17)	2.1 (0.49)	2.9 (0.16)
Fair/poor	12.3 (2.14)	23.9 (0.62)	15.9 (2.17)	21.5 (0.69)
<i>Depression medications</i>				
Yes	62.5 (9.73)	74.4 (1.42)	80.3 (5.67)	72.3 (1.73)
No	3.5 (0.61)	5.3 (0.19)	2.7 (0.47)	4.3 (0.18)

SE: standard error.

^a All p-values for chi-squared statistics comparing prevalence by immigration status were significant except for those with asterisk, which were >0.05.

Compared with the Spanish population, regardless of the survey year, immigrants were younger, less likely to be men, less educated, less likely to work in a non-manual job, more likely to have precarious employment or be unemployed, and more likely to rate their health as very good/good. Both self-reported prevalence of depression and antidepressant use were higher in the Spanish population than in the immigrant population.

The overall prevalence estimates for self-reported depression were 9.1% and 7.3% in 2014 and 2020, respectively (data not shown). In both 2014 and 2020, regardless of immigration status, the prevalence of depression was lower among men, those who were married, those with better employment status, and those who rated their health as very good/good (Table 2). Between 2014 and 2020, the prevalence of antidepressant use increased in the immigrant population (62.5% vs. 80.3%), whereas the opposite was true for native Spanish adults (74.4% in 2014 vs. 72.3% in 2020).

Table 3 shows the results of the association between immigration status and depression stratified by survey year. In the unadjusted analyses, the prevalence of depression was 52% (PR: 0.48, 95%CI: 0.36-0.65) and 31% (PR: 0.69, 95%CI: 0.53-0.88) lower among the immigrant population than among native Spanish adults in 2014 and 2020, respectively. In the adjusted analyses, these associations remained the same in 2014 and 2020. However, their strengths were attenuated for both 2014 (PR: 0.53, 95%CI: 0.50-0.79) and 2020 (PR: 0.71, 95%CI: 0.55-0.91).

We observed a similar pattern when assessing antidepressant consumption. For both survey years, immigrants reported a 69% lower use of these medications in 2014 and a 34% in 2020 lower use of this medications, as compared with their native Spanish counterparts. In the adjusted analyses, this association remained statistically significant only in 2014 (PR: 0.36, 95%CI: 0.24-0.55). We found a significant effect measure modification by immigration

Table 3

Unadjusted and adjusted prevalence ratios and 95% confidence intervals for the association of immigration status and continent of origin with depression by survey year: European Health Interview Surveys in Spain, 2014 and 2020.

	2014		2020 ^b	
	Unadjusted PR (95%CI)	Adjusted PR ^a (95%CI)	Unadjusted PR (95%CI)	Adjusted PR ^a (95%CI)
<i>Depression</i>				
Immigrant	0.48 (0.36-0.65)	0.53 (0.40-0.70)	0.69 (0.53-0.88)	0.71 (0.55-0.91)
Spanish-born	1.00	1.00	1.00	1.00
<i>Depression medications</i>				
Immigrant	0.31 (0.21-0.47)	0.36 (0.24-0.55)	0.66 (0.48-0.90)	0.74 (0.53-1.02)
Spanish-born	1.00	1.00	1.00	1.00

95%CI: 95% confidence interval; PR: prevalence ratio.

^a Adjusted for age, sex/gender, marital status, education, employment status, healthcare use, smoking status, social support, and self-rated health.

^b p-values for interactions between immigration status and survey for a) depression: 0.07, and b) depression medications: 0.01.

status and survey year for antidepressant medication use (p interaction: 0.01) but not for depression (p interaction: 0.07). However, we did not find statistically significant interactions for depression or for the use of antidepressants for immigration status with survey year and sex/gender (p interactions: 0.32 depression and 0.12 antidepressant use), or social support (p interactions: 0.28 depression and 0.31 antidepressant use).

Discussion

Our findings show a lower probability of self-reported prevalence of depression and antidepressant use among immigrants than among native Spanish adults in both EHIS 2014 and 2020. However, the association between immigration status and antidepressants was only significant in 2014 before and after adjustment. Finally, although the association of immigration status with antidepressant use differed with survey year, our findings remained the same regardless of sex/gender or social support.

Our finding of a lower prevalence of depressive symptoms and antidepressant use in the immigrant population in 2014 can be explained, at least partially, by the economic crisis that lasted from 2008 to 2014. This crisis led to the return of immigrant populations to their countries of origin between 2010 and 2015, due to economic difficulties and dramatic job losses.²⁷ Thus, it is possible that immigrants who remained had more favorable conditions in terms of social position and health status. We continue to observe immigrant health advantage in 2020 despite immigrants having more precarious employment after the economic crisis,^{28,29} being employed in positions considered precarious¹³ or essential jobs,³⁰ and having worse living conditions than their native Spanish counterparts. Moreover, other potential explanations could be the healthy immigrant effect, where there is a selection effect for healthy adults to migrate to another country, and the low use of the healthcare system observed among immigrants, especially for mental health.^{23,31,32} The latter is associated with stigma and cultural beliefs among immigrants, leading to avoiding seeking care, and thus, depression may be underestimated for this population.^{33,34} Despite the low prevalence of depression among immigrant adults, it is worth noting that the prevalence of depression in 2020 was higher among those aged 45 years or older than among their native Spanish peers. This was not the case in 2014, when a higher prevalence of depression among immigrants was observed among those aged 18–34 years. This change could be due because younger immigrants adults in 2014 encountered harsher socioeconomic conditions than their native Spanish counterparts,^{35,36} whereas older immigrants adults could have experience worst physical and mental health due to cumulative life and working conditions as well as the lock down during COVID-19 than their younger counterparts in 2020.^{37,38}

We did not find a difference in depression or antidepressant use by sex/gender or social support, regardless of migrant status

or survey year. Our findings regarding the lack of heterogeneity related to sex/gender are inconsistent with previous research suggesting that immigrant women have worse mental health than Spanish native women. However, this is not the case for immigrant men.^{8,9} There are several possible explanations for the observed differences. First, men are more likely to minimize depressive symptoms and respond differently to questions regarding their mental health conditions. Second, women are more likely to use the healthcare system,³⁹ and thus, are more likely to be aware of their mental health. Finally, women suffer from poor employment conditions and are responsible for dependent care,⁸ and their work conditions have worsened over time.⁴⁰ However, despite their access to healthcare and employment conditions, women, and especially immigrant women, coped better than men with the impact of the economic crisis,⁴¹ which could have protected their mental health. Regarding our findings related to social support, evidence suggests that social support plays a significant role in mitigating the negative association between immigration status and depression in Spain.¹⁵ However, this relationship may depend on the level of social integration and economic stability experienced by different immigrant groups, such as homeownership rates.⁴² Further research is needed to understand the specific mechanisms through which sex/gender and social support influence mental health outcomes among immigrants.

An important limitation of this study is the sample size of the immigrant population, which only allows us to analyze immigrants as a single group without accounting for key differences by country of origin or cultural background. Furthermore, we excluded immigrants from countries with HDI > 0.80 as they could have similar health status as native Spanish adults. However, we repeated the analyses including immigrants from countries with HDI > 0.80, and the results did not change significantly from the results presented in Table 3. The latter finding suggests that such exclusion was unlikely to bias our findings. Second, immigrants experiencing language barriers or precarious conditions (e.g., homelessness) were not represented in these surveys. Third, we did not account for length of stay in Spain for the immigrant population because of small size for some cells and the high correlation of the variable with immigration status. Future studies should consider length of stay, which may provide insights into the health status of the immigrant population. Fourth, we did not consider depression severity. Finally, part of the fieldwork of the 2020 survey was conducted during the state of alarm in Spain due to the COVID-19 pandemic. The latter may have precluded a greater outreach, leading to low participation in the immigrant population. Similarly, COVID-19 may have biased our results, especially for immigrants who bear a greater burden of infection and death due to their essential jobs.^{43,44} Despite these limitations, the strength of this study is that we provided a nationwide assessment of the evolution of depression prevalence among the immigrant population in Spain using the two most recent waves of the European Health Inter-

view Survey. Moreover, we observed individuals across a wide age range.

Our findings suggest that there is a protective effect for depression among the immigrant population in Spain regardless of sex/gender or social support. Future research should investigate factors contributing to depression and mental health-related outcomes in the immigrant population. The findings of such investigations could provide insights to tailor programs and interventions to address the mental health status of the population, particularly depression, regardless of immigration status, sex/gender, or social support.

What is known about the topic?

Immigrants generally exhibit a higher prevalence of depressive symptoms than native populations. In Spain, evidence for this association is mixed, with some studies reporting a higher prevalence, lower prevalence, or no difference in the prevalence of depression between immigrants and Spanish natives.

What does this study add to the literature?

Our findings suggest that there is a protective effect for depression among the immigrant population in Spain, regardless of sex/gender and social support.

What are the implications of the results?

The findings call for the need to tailor programs and interventions to address depression and mental health-related outcomes of the population regardless of immigration status, sex/gender or social support.

Availability of databases and material for replication

We used data from the National Institute of Statistics (Instituto Nacional de Estadística), Madrid, Spain, available at: https://www.ine.es/dyngs/INEbase/en/operacion.htm?c=Estadistica_C&cid=1254736176784&menu=resultados&secc=1254736195298&idp=1254735573175#tabs-1254736195298

Editor in charge

Alberto Lana.

Transparency declaration

The corresponding author, on behalf of the other authors guarantee the accuracy, transparency and honesty of the data and information contained in the study, that no relevant information has been omitted and that all discrepancies between authors have been adequately resolved and described.

Authorship contributions

L.N. Borrell conceptualized the study design, conducted the analyses, prepared the first draft of the manuscript and critically reviewed the final manuscript. N. Lanborena, E. Rodríguez-Álvarez contributed to the study conceptualization and critically reviewed the manuscript and the results. J. Díez and S. Yago-González critically reviewed the manuscript and the results. L.N. Borrell, J. Díez, S.

Yago-González, N. Lanborena and E. Rodríguez-Álvarez have read and approved the final manuscript.

Funding

This work was supported by the Multiannual Agreement reference # CM/BG/2023-002 (L.N. Borrell).

Conflicts of interest

None.

References

1. Missinne S, Bracke P. Depressive symptoms among immigrants and ethnic minorities: a population based study in 23 European countries. *Soc Psychiatry Psychiatr Epidemiol.* 2012;47:97–109.
2. Gkiouleka A, Avrami L, Kostaki A, et al. Depressive symptoms among migrants and non-migrants in Europe: documenting and explaining inequalities in times of socio-economic instability. *Eur J Public Health.* 2018;28 (Suppl 5):54–60.
3. Sempertegui GA, Baliatsas C, Knipscheer JW, Bekker MHJ. Depression among Turkish and Moroccan immigrant populations in Northwestern Europe: a systematic review of prevalence and correlates. *BMC Psychiatry.* 2023;23:402.
4. Sole-Auro A, Crimmins EM. Health of Immigrants in European countries. *Int Migr Rev.* 2008;42:861–76.
5. National Institute of Statistics of Spain. Population Figures at 1 January 2023. Available at: <https://www.ine.es/jaxiT3/Datos.htm?t=56937>.
6. Henares-Montiel J, Ruiz-Perez I, Mendoza-García O. Health inequalities between male and female immigrants in Spain after the beginning of the economic crisis. *Health Soc Care Community.* 2018;26:891–7.
7. Carrasco-Garrido P, Jimenez-García R, Barrera VH, et al. Significant differences in the use of healthcare resources of native-born and foreign born in Spain. *BMC Public Health.* 2009;9:201.
8. Gotsens M, Malmusi D, Villarroya N, et al. Health inequality between immigrants and natives in Spain: the loss of the healthy immigrant effect in times of economic crisis. *Eur J Public Health.* 2015;25:923–9.
9. Cayuela A, Malmusi D, Lopez-Jacob MJ, et al. The impact of education and socioeconomic and occupational conditions on self-perceived and mental health inequalities among immigrants and native workers in Spain. *J Immigr Minor Health.* 2015;17:1906–10.
10. Malmusi D, Palencia L, Ikram UZ, et al. Inequalities by immigrant status in depressive symptoms in Europe: the role of integration policy regimes. *Soc Psychiatry Psychiatr Epidemiol.* 2017;52:391–8.
11. Zimmerman C, Kiss L, Hossain M. Migration and health: a framework for 21st century policy-making. *PLoS Med.* 2011;8:e1001034.
12. Belvis F, Muntane F, Muntaner C, et al. What is the impact of job precariousness on depression? Risk assessment and attributable fraction in Spain. *Public Health.* 2024;231:154–7.
13. Koseoglu Ornek O, Waibel J, Wullinger P, et al. Precarious employment and migrant workers' mental health: a systematic review of quantitative and qualitative studies. *Scand J Work Environ Health.* 2022;48:327–50.
14. Gabarrell-Pascuet A, Lloret-Pineda A, Franch-Roca M, et al. Impact of perceived discrimination and coping strategies on well-being and mental health in newly-arrived migrants in Spain. *PLoS One.* 2023;18:e0294295.
15. Ortiz-Calvo E, Martínez-Ales G, Mediavilla R, et al. The role of social support and resilience in the mental health impact of the COVID-19 pandemic among healthcare workers in Spain. *J Psychiatr Res.* 2022;148:181–7.
16. Martín-Pérez A, Moreno-Fuentes FJ. Migration and citizenship law in Spain: path-dependency and policy change in a recent country of immigration. *International Migration Review.* 2018;46:625–55.
17. Huntoon L. Immigration to Spain: implications for a unified European Union immigration policy. *Int Migr Rev.* 1998;32:423–50.
18. Cabezas-Rodríguez A, Bacigalupe A, Martín U. Diagnosis and treatment of depression in Spain: are there gender inequalities? *Int J Environ Res Public Health.* 2020;17:9232.
19. United Nations Development Programme UNDP 2015. Human Development Report 2015: Work for Human Development. New York, NY. Available at: <https://hdr.undp.org/content/human-development-report-2015>.
20. Arias-de la Torre J, Vilagut G, Martín V, et al. Prevalence of major depressive disorder and association with personal and socio-economic factors. Results for Spain of the European Health Interview Survey 2014–2015. *J Affect Disord.* 2018;239:203–7.
21. Arias-de la Torre J, Vilagut G, Ronaldson A, et al. Prevalence and variability of depressive symptoms in Europe: update using representative data from the second and third waves of the European Health Interview Survey (EHIS-2 and EHIS-3). *Lancet Public Health.* 2023;8:e889–98.
22. Kocalevent RD, Berg L, Beutel ME, et al. Social support in the general population: standardization of the Oslo social support scale (OSSS-3). *BMC Psychol.* 2018;6:31.
23. Rivera B, Casal B, Currais L. The healthy immigrant effect on mental health: determinants and implications for mental health policy in Spain. *Adm Policy Ment Health.* 2016;43:616–27.

24. Backhaus I, Kino S, La Torre G, et al. Right-wing populism and self-rated health in Europe: a multilevel analysis. *J Epidemiol Community Health*. 2019;73:1116–21.
25. Cimarras-Otal C, Calderon-Larranaga A, Poblador-Plou B, et al. Association between physical activity, multimorbidity, self-rated health and functional limitation in the Spanish population. *BMC Public Health*. 2014;14:1170.
26. La Parra-Casado D, Stornes P, Solheim EF. Self-rated health and wellbeing among the working-age immigrant population in Western Europe: findings from the European Social Survey (2014) special module on the social determinants of health. *Eur J Public Health*. 2017;27 (Suppl 1):40–6.
27. Parella S, Petroff A. Migración de retorno en España: salidas de inmigrantes y programas de retorno en un contexto de crisis. *Anuario CIDOB de la Inmigración*. 2014;61–88. Available at: <https://raco.cat/index.php/AnuarioCIDOBInmigracion/article/view/288374>.
28. Jimenez-García JR, Levatino A. Stuck in a time warp? The Great Recession and the socio-occupational integration of migrants in Spain. *J Int Migr Integr*. 2023;24:1–47.
29. Arici C, Ronda-Perez E, Tamhid T, et al. Occupational health and safety of immigrant workers in Italy and Spain: a scoping review. *Int J Environ Res Public Health*. 2019;16:4416.
30. Hayward SE, Deal A, Cheng C, et al. Clinical outcomes and risk factors for COVID-19 among migrant populations in high-income countries: a systematic review. *J Migr Health*. 2021;3:100041.
31. Constant AF, Garcia-Munoz T, Neuman S, et al. A “healthy immigrant effect” or a “sick immigrant effect”? Selection and policies matter. *Eur J Health Econ*. 2018;19:103–21.
32. Elshahat S, Moffat T, Newbold KB. Understanding the healthy immigrant effect in the context of mental health challenges: a systematic critical review. *J Immigr Minor Health*. 2022;24:1564–79.
33. Nyikavaranda P, Pantelic M, Jones CJ, et al. *Int J Equity Health*. 2023;22:196.
34. Apers H, Van Praag L, Nostlinger C, et al. Interventions to improve the mental health or mental well-being of migrants and ethnic minority groups in Europe: a scoping review. *Glob Ment Health (Camb)*. 2023;10:e23.
35. Zhou J, Zhou J, Zhang H, et al. Social integration as mediator and age as moderator in social capital affecting mental health of internal migrant workers: a multi-group structural equation modeling approach. *Front Public Health*. 2022;10:865061.
36. Syed HR, Dalgard OS, Dalen I, et al. Psychosocial factors and distress: a comparison between ethnic Norwegians and ethnic Pakistanis in Oslo, Norway. *BMC Public Health*. 2006;6:182.
37. Reus-Pons M, Mulder CH, Kibele EUB, et al. Differences in the health transition patterns of migrants and non-migrants aged 50 and older in southern and western Europe (2004–2015). *BMC Med*. 2018;16:57.
38. Marin IB, Fernandez D, Ayuso-Mateos JL, et al. Healthy aging and late-life depression in Europe: does migration matter? *Front Med (Lausanne)*. 2022;9:866524.
39. Ferrara A, Grindel C, Brunori C. A longitudinal perspective to migrant health: unpacking the immigrant health paradox in Germany. *Soc Sci Med*. 2024;351:116976.
40. Fellini I. Immigrants' labour market outcomes in Italy and Spain: has the Southern European model disrupted during the crisis? *Migration Studies*. 2018;6:53–78.
41. Moreno-Colom S, Lopez-Roldan P. El impacto de la crisis en las trayectorias laborales de las mujeres inmigrantes en España. *Cuadernos de Relaciones Laborales*. 2018;36:65–87.
42. Amuedo-Dorantes C, Mundra K. Immigrant homeownership and immigration status: evidence from Spain. *Review of International Economics*. 2012;21:204–18.
43. Soriano Lopez J, Gomez Gomez JH, Ballesta-Ruiz M, et al. COVID-19, social determinants of transmission in the home. A population-based study. *Eur J Public Health*. 2024;34:427–34.
44. Perez-Muto V, Bertran MJ, Baron-Miras L, et al. Inequalities in health outcomes of SARS-CoV-2 infection by migration status in Barcelona, Spain. *Front Public Health*. 2023;11:1297025.